

Some Properties of (Non) Critical Strings

D. Kutasov*

*Joseph Henry Laboratories,
Princeton University,
Princeton, NJ 08544.*

We review some recent developments in string theory, emphasizing the importance of vacuum instabilities, their relation to the density of states, and the role of space-time fermions in non-critical string theory. We also discuss the classical dynamics of two dimensional string theory.

* Lectures at the ICTP Spring School on String Theory and Quantum Gravity, Trieste, April 1991.

1. INTRODUCTION.

In the last few years many people have been studying simple models of string theory, which correspond to string propagation in two dimensional space-time (and simple generalizations thereof). Of course, for most applications [1], [2] one needs to consider much more complicated models, however many important issues in string theory are still not understood, and the hope is that the two dimensional theory will serve as a useful toy model, in which some of these issues may be addressed. Due to the low dimension of space-time, the number of degrees of freedom in the theory is vastly smaller than in the twenty six (or ten) dimensional case. The physical on shell states include one field theoretic degree of freedom, the “tachyon” center of mass of the string (which is actually massless in two dimensions), and a discrete infinite set of massive states – the remnants of the tower of oscillator states in $D > 2$ (where the number of field theoretic degrees of freedom with mass $\leq m_0$ diverges exponentially with m_0).

Due to the absence of transverse excitations, one might expect the dynamics to be simpler in two dimensions. Indeed, following ideas of [3] [4] it has been shown [5] [6] using large N matrix model techniques, that many properties of two dimensional (2d) strings are exactly calculable to all orders in the topological (genus) expansion. The simplicity of these models is closely related to an underlying free fermion structure [3] – [7].

There are two main potential domains of application of the results of [5], [6]. One is two dimensional (world sheet) quantum gravity. The implications of matrix models for quantum gravity have been recently discussed in [8], [9], [10]. The second application, which will be of main interest to us here, is to critical (unified) string theory. The two are of course closely connected, and we will mention some aspects of this relation as we go along. The approach we will take is to try and understand the matrix model results in the continuum path integral formalism [1], and use this understanding to shed light on the structure of the theory, in particular its space-time dynamics. This program is far from complete; we will review what has been achieved and discuss some of the open problems.

We will take a broader point of view, presenting 2d strings in the context of higher dimensional string theory, which is ultimately what we’re interested in. This will also help to try and identify features of general validity, and those that are special to two dimensions. With that in mind, we will also investigate some string models in two dimensions which are more difficult to treat (or in some cases even formulate) using matrix models.

In the continuum formulation of 2d quantum gravity [11] [12], we are given a general conformal field theory (CFT) with action $\mathcal{S}_M(g)$ (on a Riemann surface with metric g_{ab}),

central charge c_M , and we have to integrate over all metrics g on manifolds of fixed topology (genus). In the conformal gauge $g_{ab} = e^\phi \hat{g}_{ab}$, the dynamical metric g is represented by the Liouville mode ϕ ; the action is:

$$\mathcal{S} = \mathcal{S}_L(\hat{g}) + \mathcal{S}_M(\hat{g}) \quad (1.1)$$

where

$$\mathcal{S}_L(\hat{g}) = \frac{1}{2\pi} \int \sqrt{\hat{g}} [\hat{g}^{ab} \partial_a \phi \partial_b \phi - \frac{Q}{4} \hat{R} \phi + 2\mu e^{\alpha_+ \phi}] \quad (1.2)$$

Q and α_+ are parameters which are fixed by gauge invariance [13]. It is very useful to think about the Liouville mode as a target space coordinate, so that (1.1) is a particular background of the critical string system in two dimensions [14]. Therefore, we have to set the total central charge of matter (c_M) and Liouville ($c_L = 1 + 3Q^2$) to 26:

$$Q = \sqrt{\frac{25 - c_M}{3}} \quad (1.3)$$

Spinless matter primary operators of dimension $\Delta (= \bar{\Delta})$ V_Δ acquire Liouville dependence,

$$T_\Delta(\phi) = V_\Delta \exp(\beta_\Delta \phi) \quad (1.4)$$

with

$$\Delta - \frac{1}{2}\beta(\beta + Q) = 1 \quad (1.5)$$

In particular, putting $\Delta = 0$ in (1.5) fixes α_+ in (1.2). As usual in critical string theory, (1.5) is the mass shell condition (or the linearized equation of motion for small excitations). The vertex operators T_Δ are related to the wave functions of the corresponding states Ψ_Δ by $T_\Delta(\phi) = g_{\text{st}}(\phi) \Psi_\Delta(\phi)$, where unlike the “usual” cases [1], the string coupling g_{st} is *not* constant in general (1.2); rather we have $g_{\text{st}} = g_0 \exp(-\frac{Q}{2}\phi)$. Therefore,

$$\Psi_\Delta = V_\Delta e^{(\beta + \frac{Q}{2})\phi} \quad (1.6)$$

and $E = \beta + \frac{Q}{2}$ is identified as the Liouville momentum of the state described by Ψ . The form (1.6) is actually only an asymptotic approximation of the exact wave function. It is valid in the region $e^{\alpha_+ \phi} \ll 1$ ($\phi \rightarrow \infty$ in our conventions). The exact wave function in two dimensional string theory is known [9] to satisfy the Wheeler de Witt equation,

$$\left[-\frac{\partial^2}{\partial \phi^2} + \mu \exp(\alpha_+ \phi) + \nu^2 \right] \Psi_\Delta(\phi) = 0 \quad (1.7)$$

where $\nu^2 = 2\Delta - \frac{c_M - 1}{12}$. Eq. (1.7) was derived [9] in the matrix model approach. In the continuum formalism, (1.7) holds in a minisuperspace approximation; it was not yet derived in the full theory. It is very natural to interpret ϕ as Euclidean time. Eq. (1.5) takes then the form:

$$E^2 = m^2; \quad m^2 = 2\Delta - \frac{c_M - 1}{12} \quad (1.8)$$

The wave function Ψ_Δ with energy E describes one quantum mechanical degree of freedom in space-time. If, as in the minimal models of [15], the matter system contains a finite number of spinless primaries V_Δ , the corresponding string theory reduces to quantum mechanics of a finite number of degrees of freedom. In general, of course, the number of matter primaries is enormous, and the number of degrees of freedom is much larger than in field theory. If there are some non compact dimensions X_i , such that $V_\Delta(X) = e^{ik \cdot X} \tilde{V}_h$, $\Delta = \frac{1}{2}k^2 + h$, we can rewrite (1.8) as

$$E^2 = k^2 + m^2, \quad m^2 = 2h - \frac{c_M - 1}{12}$$

and V describes one field theoretic degree of freedom in the appropriate dimension.

From (1.8) we see that operators with $\Delta < \frac{c_M - 1}{24}$ in the matter CFT correspond in space-time to on shell tachyons, which lead (as in field theory) to IR instabilities. In field theory, these IR divergences imply that we are expanding around the wrong vacuum, and should move to a new, stable one. We will see that in string theory, as it is currently understood, the situation is surprisingly different.

From the point of view of 2d gravity, tachyons correspond to normalizable wave functions (1.6), while states with positive m^2 are described by wave functions which are exponentially supported at $\phi \rightarrow \pm\infty$ (and are of course non normalizable). The world sheet interpretation of this [8] is that normalizable (tachyonic) wave functions are supported mainly on surfaces with finite size holes *in the dynamical metric* g_{ab} ; they describe macroscopic states, and if we perturb the action (1.1) by the corresponding operators, the dynamical surface is destabilized by the multitude of holes that are created. Non normalizable wave functions with $E > 0$ are supported on very small surfaces in the dynamical metric, and therefore describe small disturbances of the surface (microscopic states). The corresponding operators can be (and were) studied in the matrix model approach. Wave functions with $E < 0$ describe very large disturbances of the surface, and it is not known how to treat them in the matrix model (although there are some plausible suggestions).

Thus, there is a nice correspondence between IR instabilities in space time (tachyons) and on the world sheet (macroscopic states which create holes in the surface). The role of states with $E < 0$ is not well understood at this time, and is in fact one of the important remaining problems, which is related to many properties of the theory, as we will see below. Note that by (1.8) the familiar relation between tachyons and relevant matter operators is only true for $c_M = 25$. In general, for $c_M < 25$ relevant operators with $\frac{c_M-1}{24} < \Delta < 1$ do not create instabilities; for $c_M > 25$ we have the unintuitive situation that irrelevant matter operators with $1 < \Delta < \frac{c_M-1}{24}$ are tachyonic. The region $c_M > 25$ has been less studied than $c_M < 25$, and we will not discuss it further here.

For tachyonic operators, β in (1.4) is complex (1.8). In the original work of [12], [13] (see also [16] [17]) it was noted that the identity operator ($V_\Delta = 1$ in (1.4)) becomes tachyonic when $c_M > 1$. Since the gravitational coupling constant is proportional to $\frac{1}{26-c_M}$ [11], it has been suggested that at $c_M = 1$ a certain transition between “weak gravity” $c_M < 1$ and “strong gravity” $c_M > 1$ takes place. Taken naively, this point of view was not completely satisfactory, since it is easy to construct matter theories where $c_M < 1$, but there exist tachyonic operators in the spectrum; it is not likely that the identity operator plays a special role in this context. The issue was later clarified in [8], [18]; the current understanding is that the crucial quantity is not the gravitational coupling constant, but rather the *density of states* of the theory. Gravity has $\frac{1}{2}n(n-3)$ degrees of freedom in n dimensions; for $n = 2$ this is -1 . If the matter theory has more than “one degree of freedom”, in a sense which we will make precise below, the full string theory contains tachyons. Therefore, the invariant way to describe the famous “ $c = 1$ barrier” of [12], is that *2d gravity coupled to matter with more than one field theoretic degree of freedom is unstable*. The relation between the existence of tachyonic excitations around a certain vacuum, and the number of states in it implies that if it is possible at all to turn on expectation values of the various fields and move from a tachyonic vacuum to a stable one in string theory, the new stable vacuum is always “trivial” (at least in bosonic string theory).

Hence, the density of states is of major importance in string theory, as are tachyonic instabilities. Section 2 of these notes is devoted to a precise definition of the density of states and its relation to the presence of tachyons and stability. We also discuss the implications of this relation for string dynamics.

Since the only stable vacua of bosonic string theory (where by bosonic we mean vacua with only bosonic space-time excitations; e.g. the fermionic string falls into this category

as well) are those which are essentially two dimensional field theories in space-time, we are led in section 3 to consider theories with space-time fermions [19]. We construct explicitly a large set of theories with a number of degrees of freedom varying between that of a two dimensional field theory and that of the ten dimensional superstring, all of which are stable solutions of the equations of motion of string theory. This is possible since space time fermions contribute a negative amount to the “number of degrees of freedom”, and a theory with many bosons and many fermions can still have a vanishingly small total “density of states”. This number is required to be small for stability. The theories thus obtained have many of the favorable properties of critical superstring models, but exhibit some puzzling features as well (such as continuous breaking of SUSY), mostly related to their time dependence.

In section 4 we describe the known facts about classical dynamics in two dimensional string theory. One of the issues of interest is the exact form of the classical equations of motion of the string. The usual way one obtains those in string theory is by studying scattering amplitudes of on shell fields. Calculating the correlation functions of T_Δ (1.4) involves solving the interacting Liouville theory (1.2). Despite many efforts [16], [17], [8], [20], this is still an open problem. What saves the day is that there exists a class of amplitudes which contain most of the information, and are simply calculable. To understand this, it is useful to focus on the dynamics of the zero mode of the Liouville field ϕ (the minisuperspace approximation in 2d gravity). The main source of the complications in Liouville theory is the “cosmological” interaction in (1.2). From the space-time point of view, this corresponds to a (zero X momentum) tachyon condensate, while on the world sheet this is a potential for ϕ . The reason why it is needed is to keep the Liouville field away from the region $\phi \rightarrow -\infty$ where the string coupling $g_{st} \rightarrow \infty$. Many amplitudes diverge when $\mu \rightarrow 0$ in (1.2). By KPZ scaling [12], the amplitudes $\langle T_{\Delta_1} \dots T_{\Delta_n} \rangle$ are proportional to μ^s , where

$$\sum_{i=1}^N \beta_i + \alpha_+ s = -Q \quad (1.9)$$

In general, all amplitudes are non zero – there is no Liouville momentum conservation (1.9). Instead, the ϕ zero mode integral, which can be explicitly performed, yields [21]:

$$\langle T_{\Delta_1} \dots T_{\Delta_n} \rangle = \left(\frac{\mu}{\pi} \right)^s \Gamma(-s) \langle T_{\Delta_1} \dots T_{\Delta_n} \left[\int \exp(\alpha_+ \phi) \right]^s \rangle_{\mu=0} \quad (1.10)$$

where the correlator on the r.h.s. is understood to exclude the Liouville zero mode. One can see from (1.10) that the amplitudes with $s = 0$ are special. The ϕ zero mode integral is given for them by:

$$\int_{-\infty}^{\infty} d\phi_0 \exp(-\mu e^{\alpha_+ \phi_0})$$

This diverges from $\phi \rightarrow \infty$; imposing a UV cutoff $\phi_0 \leq \phi_{UV}$, we find $\log \mu$ multiplied by a free field amplitude for the Liouville field (i.e. Liouville is treated as a Feigin-Fuchs field, and the cosmological interaction is absent). Of course, we still have to multiply by the matter contribution and integrate over moduli, but this reduces the calculation to well defined integrals; similar representations for generic Liouville amplitudes (1.10) are not known.

Why does this simplification of the amplitudes with $s = 0$ occur? A major clue is provided by the observation that these amplitudes are insensitive to the particular form of the “wall” that keeps the Liouville field away from $\phi = -\infty$. For example, replacing $e^{\alpha_+ \phi}$ by a general T_Δ would do the job, and while generic amplitudes depend strongly on the choice of the condensate T_Δ , the $s = 0$ ones do not (apart from trivial overall factors). The existence of the Liouville wall is of course a boundary effect, hence the $s = 0$ amplitudes correspond to *bulk scattering*, which is clearly independent of the form of the boundary at strong coupling. The reason for the simplicity is that in the bulk of the ϕ volume, all the complications of Liouville theory are irrelevant, and the Liouville field is free. The overall factor of $\log \frac{1}{\mu}$ which appears in all bulk amplitudes is interpreted as the volume of the Liouville coordinate, as appropriate for bulk effects.

A similar simplification occurs in (1.10) for all $s \in Z_+$. The power of $\int \exp(\alpha_+ \phi)$ on the r.h.s. becomes then a positive integer; the divergence of $\mu^s \Gamma(-s)$ is interpreted again as¹: $\mu^s \Gamma(-s) \simeq \mu^s \log \mu$. The space-time interpretation is as before: now it involves bulk interaction of the scattering particles $T_{\Delta_1} \dots T_{\Delta_n}$, with the zero momentum tachyons that form the Liouville wall.

The bulk amplitudes provide us with the required probe of the dynamics of D dimensional string theories; they can be defined and evaluated for all string vacua (all matter

¹ This can be obtained by starting with the fixed area amplitude, using $\mu^s \Gamma(-s) = \int_0^\infty dA A^{-s-1} e^{-\mu A}$. The divergence at $s \in Z_+$ is interpreted then as a small area divergence in the integral over areas, which can be regulated by a small area (UV) cutoff, as before, yielding the stated result.

CFT's M), while general Liouville amplitudes (1.10) are expected to be much more subtle for $D > 2$. The bulk amplitudes should also be sufficient to deduce the equations of motion of the theory, and compare the dynamics around different vacua. In principle one can construct from them a space-time Lagrangian which gives those amplitudes, and then incorporate boundary effects to deduce all the amplitudes of the theory [22].

In view of the above discussion, section 4 is devoted to the bulk tree level S – matrix in two dimensional string theory. Surprisingly, the moduli integrals can be explicitly evaluated in this case, essentially due to the same simplification of the dynamics of the theory, as that leading to the free fermion structure of the corresponding matrix models. In the continuum formalism, the simplicity is due to a partial decoupling of a certain infinite set of discrete string states. We review the results and mention some problems still left in the continuum approach to 2d strings. We also compare the situation in 2d closed string theory to that found in open 2d string theory. Section 5 contains some concluding remarks.

2. DENSITY OF STATES AND TACHYONS IN STRING THEORY.

In this section we will show that the density of states (or number of degrees of freedom) of the system plays a central role in two dimensional quantum gravity and string theory. In particular, we will show that matter with ‘too many’ degrees of freedom can not be consistently coupled to gravity (under mild assumptions) due to the appearance of tachyons. We believe that this restriction on the number of states in quantum mechanical generally covariant theories is more general.

We start by defining the density of states. The most general situation we are interested in is an arbitrary vacuum of critical string theory, i.e. an arbitrary conformal field theory with $c = 26$ (or $\hat{c} = 10$). As we saw in section 1, gravity coupled to a matter CFT with any c_M is a particular example, with the missing central charge $26 - c_M$ supplied by the Liouville CFT. We’ll start by considering this special case, and then generalize. To count states it is convenient to evaluate the torus partition sum:

$$Z(\tau) = \text{Tr} q^{L_0 - \frac{c}{24}} \bar{q}^{\bar{L}_0 - \frac{\bar{c}}{24}} \quad (2.1)$$

where $q = e^{2\pi i\tau}$, and $\tau = \tau_1 + i\tau_2$ is the complex modulus of the torus.

The string partition sum (2.1) factorizes into a product of three contributions, from ghosts (for which a factor of $(-)^G$, G – ghost number, must be inserted in (2.1)), Liouville and the “matter” CFT. The ghosts contribute to $Z(\tau)$ a factor of $|\eta(\tau)|^4$. The Liouville

contribution is that of a free scalar field [21] $\log \mu \left(\sqrt{\tau_2} |\eta(\tau)|^2 \right)^{-1}$; the cosmological term in (1.2) enters trivially, for the same reasons as in section 1 – the one loop free energy is a bulk amplitude; hence it is proportional to the volume $\log \mu$, with the coefficient given by a free field amplitude. Finally, the matter is described by a partition sum $Z_{\text{CFT}}(\tau)$ (defined as in (2.1)).

The total partition sum is (after multiplying by a factor of τ_2 for later convenience)

$$Z_{\text{string}} = \sqrt{\tau_2} |\eta|^2 Z_{\text{CFT}} \quad (2.2)$$

Modular invariance of $Z_{\text{string}}(\tau)$ is a fundamental principle in string theory. We will assume it throughout our discussion. In ordinary CFT, the partition sum (2.1) can be used to count states in the theory. One considers the behavior of the partition sum (Z_{CFT} or Z_{string}), when $\tau_1 = 0, \tau_2 = \beta \rightarrow 0$, where $Z(\beta) = \text{Tr } e^{-\beta E}$ has the form:

$$Z(\beta) \sim \beta^r \exp\left(\frac{\pi c_{\text{eff}}}{6\beta}\right) \quad (2.3)$$

Clearly, c_{eff} is a measure of the number of degrees of freedom of the theory, since in the limit $\beta \rightarrow 0$ all states contribute 1 to $Z(\beta)$ (2.1). By modular invariance $Z(\beta) = Z(\frac{1}{\beta})$, and (2.1), c_{eff} is related to the lowest lying states in the spectrum:

$$\begin{aligned} c_{\text{eff}} &= c - 24\Delta_m \\ \Delta_m &= \frac{1}{2} \min(\Delta + \bar{\Delta}) \end{aligned} \quad (2.4)$$

In unitary CFT $\Delta_m = 0$, and $c_{\text{eff}} = c$. So the central charge measures the number of degrees of freedom [23]. In non unitary theories, Δ_m is generically negative and $c_{\text{eff}} > c$. In Liouville theory, $c_{\text{eff}} = 1$ ($\Delta_m > 0$). Thus $c(= 1 + 3Q^2)$ is not a good measure of the density of states, which is that of a free scalar field for all Q .

In string theory, c_{eff} is not a good measure of the density of states since not all states contributing to (2.3) are physical – we have to impose the constraint $L_0 = \bar{L}_0$. This is implemented by integrating over τ_1 ; we define:

$$G(\tau_2) = \int_{-\frac{1}{2}}^{\frac{1}{2}} d\tau_1 Z_{\text{string}} \quad (2.5)$$

By modular invariance we know that $Z_{\text{string}}(\tau_1 + 1, \tau_2) = Z_{\text{string}}(\tau_1, \tau_2)$, hence only states of integer spin contribute to Z_{string} ($\Delta - \bar{\Delta} \in \mathbb{Z}$). Transforming to $G(\tau_2)$ (2.5) eliminates

contributions of all states except those with $\Delta = \bar{\Delta}$. The function $G(\tau_2)$ has a simple space-time interpretation. It is related to the one loop free energy of the particle excitations of the string: $\text{Tr}(-)^F \log(p^2 + m^2)$ (F is the space-time fermion number). In a proper time representation, the one loop free energy Ω is given by:

$$\Omega = \text{Tr}(-)^F \log(p^2 + m^2) = \int_0^\infty \frac{ds}{s} \sum_n (-1)^{F_n} \int d^D p e^{-s(p^2 + m_n^2)} \quad (2.6)$$

where p is the momentum in the non compact directions and the sum over n runs over the space-time excitations. The function $G(\tau_2)$ of (2.5) is related to the integrand in (2.6):

$$G\left(\frac{s}{2\pi}\right) = s \sum_n (-1)^{F_n} \int d^D p e^{-s(p^2 + m_n^2)} \quad (2.7)$$

One consequence of this is that all states should contribute positive amounts to $G(\tau_2)$, unless there are space-time fermions in the spectrum. From the definition of $G(\tau_2)$ (2.2), (2.5), this is far from clear. While the CFT partition sum as well as that of Liouville satisfy this property, being traces over Hilbert spaces with positive weights, ghost oscillators flip the sign of the contributions to (2.1) (due to the factor of $(-)^G$ in the definition of the ghost partition sum). Therefore, even in bosonic string theory there may appear physical states at non trivial ghost numbers, which contribute a negative amount to $G(\tau_2)$ (2.5). Such states indeed do appear at discrete values of the momenta p . They seem to decouple from the dynamics due to the ghost numbers, and their role is not completely clear. In more than two dimensions the statistical weight of these states is low and they can be ignored (except perhaps as generators of symmetries). In two dimensional space-time they seem to be closely related to the symmetry structure of the theory [24]. In any event, the field theoretic degrees of freedom, over which the sum in (2.7) runs, occur at zero ghost number and contribute with positive sign to the partition sum.

A state with $\Delta = \bar{\Delta}$ in the CFT contributes to $G(\tau_2)$, an amount $e^{-4\pi\tau_2(\Delta - \frac{c_M-1}{24})}$. If $\Delta < \frac{c_M-1}{24}$, the Schwinger integral (2.6) develops an IR divergence (from $\frac{s}{2\pi} = \tau_2 \rightarrow \infty$). This IR divergence is due to the tachyon instability. Hence, the behavior of $G(\tau_2)$ as $\tau_2 \rightarrow \infty$ probes the existence of tachyons in the spectrum. On the other hand, its behavior as $\tau_2 \rightarrow 0$ is a measure of the density of states of the theory, since in that limit all states contribute 1 to G . Now, the theories we are describing here all have an infinite number of states, therefore generically G diverges as $\tau_2 \rightarrow 0$; however we can estimate the density of states by measuring how fast G diverges in that limit.

On general grounds we know that as $\tau_2 \rightarrow 0$, $G(\tau_2) \simeq \tau_2^x e^{\frac{y}{\tau_2}}$. Therefore it is natural to define:

$$c_{\text{string}} - 1 = \lim_{\tau_2 \rightarrow 0} \frac{6\tau_2}{\pi} \log G(\tau_2) \quad (2.8)$$

as a measure of the number of degrees of freedom. The -1 on the left hand side of (2.8) reflects the contribution of the gravity sector to the density of states mentioned above. If one takes, e.g., the matter CFT to be one scalar field or a minimal model [15], it is easy to see that $c_{\text{string}} = 1$. Thus the minimal models and two dimensional string theory have the same density of states. In general, $c_{\text{string}} > 1$ for theories without space-time fermions. We will see soon that such theories always contain tachyons.

As explained above, tachyons cause an IR divergence in Ω , thus finiteness of Ω is a good measure of the existence of tachyons (in fermionic string theories this is strictly true only for theories without tachyonic space-time fermions, e.g. theories with unitary matter). Of course, the field theoretic expression for Ω (2.6) has in addition to the IR tachyon divergence which we are interested in, a UV divergence (from $s = 0$). This is usually dealt with by a proper time cutoff: $\int_0^\infty ds \rightarrow \int_{\Lambda^{-2}}^\infty ds$. String theory removes this divergence by a different mechanism [1]. One notes that by (2.5), (2.6), (2.7), $\Omega = \int \frac{d^2\tau}{\tau_2^2} Z_{\text{string}}(\tau)$, where the integral runs over the half infinite strip $\tau_2 \geq 0$, $|\tau_1| \leq \frac{1}{2}$. This may diverge due to modular invariance of Z_{string} since the strip contains an infinite number of copies of a fundamental domain of the modular group. If this is the only source of divergence, one can cure it by restricting the integral to a fundamental domain $\mathcal{F}$ of the modular group, $|\tau| \geq 1$, thus avoiding the “UV region” $\tau_2 \rightarrow 0$. But this “stringy regularization” actually teaches us something very interesting about the theory. Suppose there are no tachyons in the spectrum. Then the integral $\int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^2} Z_{\text{string}}$ is finite, and the only possible source of divergence of the integral over the strip is the volume of the modular group. We can try to estimate this divergence by using the “field theoretic” cutoff: integrate $\int \frac{d^2\tau}{\tau_2^2}$ over the cutoff strip: $|\tau_1| \leq \frac{1}{2}$, $\tau_2 > \frac{1}{\Lambda^2}$. The integral of Z_{string} over the cutoff strip should diverge at the same rate as the (regularized) volume of the modular group, which is given by $\int_{\Lambda^{-2}}^\infty \frac{d\tau_2}{\tau_2^2} \int_{-\frac{1}{2}}^{\frac{1}{2}} d\tau_1 1 = \Lambda^2$. Hence we learn that from the space-time point of view, tachyon free string theories have a free energy $\Omega(\Lambda)$, which diverges (at most) as $\Omega \simeq \Lambda^2$ as we remove the cutoff. But in space-time the rate of divergence of Ω measures the density of states of the theory. The behavior we find in tachyon free string theory indicates a very small number of states. Even a theory with one bosonic field in D dimensions would have $\Omega(\Lambda) \simeq \Lambda^D$. In string theory we generically have an infinite number of space-time fields,

so the behavior we find is even more unusual. We would expect $\Lambda^a e^{b\Lambda}$ in general, due to the Hagedorn-like growth of the number of states with mass. By the above arguments, such theories are always tachyonic. The only theories that can be tachyon free are those that exhibit the general features of 2D field theories!

Pushing these ideas one step further, we can derive a more quantitative correspondence between Ω and the number of states. If the divergence of (2.6) is indeed due to the volume of the modular group, we should have a relation of the form:

$$\lim_{\Lambda \rightarrow \infty} \frac{\int_{\Lambda^{-2}}^{\infty} \frac{d\tau_2}{\tau_2^2} \int_{-\frac{1}{2}}^{\frac{1}{2}} d\tau_1 Z_{\text{string}}}{\int_{\Lambda^{-2}}^{\infty} \frac{d\tau_2}{\tau_2^2} \int_{-\frac{1}{2}}^{\frac{1}{2}} d\tau_1 1} = \frac{\int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^2} Z_{\text{string}}}{\int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^2} 1} \quad (2.9)$$

Equation (2.9) is the statement that the two regularizations of Ω using the field theoretic proper time cutoff, and the stringy modular invariant cutoff, are equivalent. This is in fact only true under certain assumptions, which we will soon state, but assuming it is true, we derive by evaluating the l.h.s. the following relation:

$$\int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^2} Z_{\text{string}} = \lim_{\Lambda \rightarrow \infty} \frac{\pi}{3\Lambda^2} \int_{\Lambda^{-2}}^{\infty} \frac{d\tau_2}{\tau_2^2} G(\tau_2) = \lim_{\Lambda \rightarrow \infty} \frac{\pi}{3} G(\Lambda^{-2}) \quad (2.10)$$

This relation states that the (regularized) one loop free energy is equal (up to a constant) to the number of states of the theory $G(0)$. Before going on to prove (2.10), we would like to make several comments on its significance and implications.

Eq. (2.10) is a statement about modular invariant functions (obeying certain conditions to be stated below). We have presented it for the case of a CFT coupled to Liouville, however the discussion applies to all string vacua which are described by modular invariant partition sums. Important generalizations include fermionic strings, superstrings, and heterotic strings, as well as arbitrary CFT's with $c = 26$.

The l.h.s. of (2.10) diverges iff there are tachyons in the spectrum. Hence, if there are no tachyons, the r.h.s. must also be finite. Looking back at (2.7), we see that this implies that the theory has the number of states of a two dimensional field theory with a finite number of fields. If there are no space-time fermions, this means that the theory contains far fewer states than generic string theories [1]. The only bosonic string theories without tachyons are two dimensional (e.g. the $c = 1$ or minimal models coupled to gravity, and the coset model of [25]). This means that the role of tachyons in string theory is more fundamental than in field theory. Unlike there, it doesn't seem likely that tachyons can be gotten rid of by shifting to a nearby vacuum. For that to happen, one of two scenarios

must occur: either space-time fermions must somehow dynamically appear, or the new stable vacuum must effectively be a two dimensional string theory. Starting e.g. from the 26 dimensional bosonic string, neither possibility seems likely. In general, it seems unlikely that turning on expectation values of fields can change c_{string} , which should be invariant under small perturbations. The fate of string vacua containing tachyons (or 2D quantum gravity systems with “too much” matter) remains unclear.

Fermionic string theories with a non chiral GSO projection contain also only bosonic excitations in space-time. Hence the situation there is completely analogous to the bosonic case – tachyon free theories have very few states (the number of states is again as in two dimensional field theory). Theories with many states contain tachyons. It is well known that non trivial theories exist in this case, but those involve a *chiral* GSO projection (e.g. the ten dimensional superstring [1]). It is important to note that such theories are not necessarily space-time supersymmetric (one example of a non supersymmetric theory is the $O(16) \times O(16)$ theory [1]). However, it is known that they always contain space-time fermions. Our discussion uncovers a surprising feature of these theories: even though space-time SUSY is absent in general, tachyon free superstrings are *asymptotically supersymmetric*. In other words, while the number of bosons and fermions is not the same energy level by energy level, at high enough energy, the total numbers of bosons and fermions up to that energy are the same to fantastic precision. More precisely, in this case it is natural to write $G(s)$ (2.7) as $G(s) = G_B(s) - G_F(s)$, where G_B and G_F measure the contributions of bosons and fermions to the free energy (2.7), and the relative minus sign is due to spin statistics; then in general we have $G_B(s), G_F(s) \simeq s^a e^{\frac{b}{s}}$, while $G_B(s) - G_F(s) \rightarrow \text{const}$ as $s \rightarrow 0$ (again, by using (2.10) and finiteness of its l.h.s.). Hence bosons and fermions almost cancel in any tachyon free string theory. This “asymptotic supersymmetry” is a very puzzling phenomenon, whose full implications are still not understood. Non supersymmetric tachyon free superstring theories resemble models with spontaneously broken SUSY (of course the scale of breaking is the Planck scale in general). Whether this is more than a formal similarity remains to be seen. Note also that the one loop cosmological constant in tachyon free string theories is unnaturally small from the space-time point of view (although still much too large). From our point of view, Ω can vanish without space-time supersymmetry. This requires only that G_B and G_F cancel precisely as $s \rightarrow 0$. We don’t know whether or why this should be the case.

In theories with space-time fermions, one may also define $\tilde{G}(s) = G_B(s) + G_F(s)$, which counts the total density of states of bosons *plus* fermions. This quantity behaves

when $s \rightarrow 0$ as: $\tilde{G}(s) \simeq s^a e^{\frac{b}{s}}$, and determines the thermal properties (e.g. the Hagedorn temperature) of the theory. It is not clear whether models with $b > 0$ (such as those of section 3) exhibit any simplifying features due to the fact that $G_B(s) - G_F(s) \simeq 0$ (as $s \rightarrow 0$).

Our remaining task in this section is to state precisely and prove (2.10).

Theorem: Let $Z_{\text{string}}(\tau)$ be a modular invariant function, which is finite throughout the fundamental domain $\mathcal{F}$, except perhaps at $\tau_2 = \infty$, such that:

- 1) $\int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^2} Z_{\text{string}} = \text{finite}$. It is implied in the definition of the above integral that for $\tau_2 \gg 1$ we first perform the integral over $-\frac{1}{2} \leq \tau_1 \leq \frac{1}{2}$, and then integrate over τ_2 .
- 2) As $\tau_2 \rightarrow \infty$, $Z_{\text{string}} \simeq \tau_2^x q^a \bar{q}^b$ ($q = e^{2\pi i \tau}$) where $a, b > -1$.

Then:

$$\int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^2} Z_{\text{string}} = \frac{\pi}{3} \lim_{\tau_2 \rightarrow 0} \int_{-\frac{1}{2}}^{\frac{1}{2}} d\tau_1 Z_{\text{string}}(\tau_1, \tau_2) \quad (2.11)$$

Proof: Consider the function:

$$F(R) = \int_0^\infty \frac{d\tau_2}{\tau_2^2} \int_{-\frac{1}{2}}^{\frac{1}{2}} d\tau_1 Z_{\text{string}}(\tau) \sum_{r \neq 0}^\infty \exp\left(\frac{-\pi R^2 r^2}{\tau_2}\right) \quad (2.12)$$

This is a cutoff version of (2.6), with a “modular invariant” cutoff. R plays the role of Λ^{-1} in the discussion above. Our purpose is to relate the integral over the strip $F(R)$ (2.12) to an integral over the fundamental domain $\mathcal{F}$. Naively, this can be done using [26]:

$$\int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^2} Z_{\text{string}}(\tau) \exp\left(-\pi R^2 \frac{|n - m\tau|^2}{\tau_2}\right) = \int_{\alpha \cdot \mathcal{F}} \frac{d^2\tau}{\tau_2^2} Z_{\text{string}}(\tau) \exp\left(-\pi R^2 \frac{r^2}{\tau_2}\right) \quad (2.13)$$

where α is the modular transformation taking $e^{-\pi R^2 \frac{|n - m\tau|^2}{\tau_2}} \rightarrow e^{-\pi R^2 \frac{r^2}{\tau_2}}$. We can write the sum over r and integral over the strip (2.12) as a sum over r, α and an integral over the fundamental domain $\mathcal{F}$ (since the sum over $\alpha \cdot \mathcal{F}$ generates the strip), and then replace the sum over r, α by a sum over n, m (2.13) (and integral over $\mathcal{F}$). This would suggest:

$$\int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^2} Z_{\text{string}}(\tau) \sum_{n, m \in \mathbb{Z}} \exp\left(-\pi R^2 \frac{|n - m\tau|^2}{\tau_2}\right) = F(R) + \int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^2} Z_{\text{string}} \quad (2.14)$$

This is unfortunately too naive. The problem is that in many interesting situations (most notably heterotic strings), the theory contains “unphysical tachyons” (with $L_0 \neq \bar{L}_0$), which lead to divergent terms as $q \rightarrow 0$, but disappear after the τ_1 integration. In such situations the above integrals (e.g. (2.13)) are not absolutely convergent. We have to

integrate the l.h.s. of (2.13) over τ_1 first. In such cases, the simple order of integration on $\mathcal{F}$ translates to a complicated prescription on $\alpha \cdot \mathcal{F}$, different for each modular transformation α . Since the integral depends on the order of integration, (2.14) is in general invalid. Clearly, in general the situation is out of control and the argument of [26] can only be applied if we're dealing with absolutely convergent integrals. Fortunately, this is the case if R in (2.14) is large enough. By condition (2) of the theorem, we are dealing with $Z_{\text{string}}(\tau)$ such that there exists a R_0 such that for all $R > R_0$ the l.h.s. of (2.13) is independent of the order of integration, as long as $m \neq 0$. But, since we use (2.13) only for terms with $m \neq 0$ (otherwise $\alpha = 1$ in (2.13)), for $R > R_0$, (2.14) is valid. Now we can continue (2.14) analytically to $R < R_0$ and it must still hold. The reason is that both the l.h.s. and the r.h.s. of (2.14) define analytic functions of R in a strip $\text{Im}R < \epsilon$, $R > 0$. For the l.h.s. this is clear, since the only divergences come from $\tau_2 \rightarrow \infty$, but by condition (2) of the theorem no such divergence is possible for the function and all its derivatives w.r.t. R . The r.h.s. is analytic since it is equal to the l.h.s. for $R > R_0$, and the analytic continuation is unique. We conclude that (2.14) is correct all the way down to $R = 0$. Poisson resummation of the l.h.s. gives

$$\frac{1}{R} \int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^{\frac{3}{2}}} Z_{\text{string}}(\tau) \sum_{n,m \in \mathbb{Z}} q^{\frac{1}{4}(\frac{n}{R} + mR)^2} \bar{q}^{\frac{1}{4}(\frac{n}{R} - mR)^2} = F(R) + \int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^2} Z_{\text{string}}(\tau) \quad (2.15)$$

As $R \rightarrow 0$ both sides behave as $\frac{1}{R^2}$. Equating the coefficients gives (2.11).

The conditions of the theorem are very natural. Condition 1 is obvious; the main reason to require condition 2 is that it is equivalent to unitarity when $c < c_{\text{critical}}$. E.g. in (2.2) we have $Z_{\text{string}} \simeq q^{\Delta - \frac{c_M - 1}{24}} \bar{q}^{\bar{\Delta} - \frac{c_M - 1}{24}}$ (as $q \rightarrow 0$), and as long as $c_M < 25$, if all $\Delta \geq 0$ (as is the case in unitary CFT's), condition 2 is satisfied. The reason one expects the theorem to break down in general for non unitary matter, is that then the Ramond sector may contain tachyons as well as the NS sector. But in that case, bosonic and fermionic tachyons may cancel in the expression for the free energy, giving rise to a finite l.h.s. of (2.11), but still contribute to a large density of states, such that the r.h.s. of (2.11) is infinite. For $c > c_{\text{critical}}$, even unitary theories may violate the theorem. This region is not understood, as mentioned above. Some subtleties may also occur on the boundary between the two regions – the critical (heterotic) string (see [18] for details).

3. NON – CRITICAL SUPERSTRINGS.

In section 2 we saw that all non trivial theories of strings without tachyon instabilities must contain space-time fermions, in fact essentially the same number of fermions and of

bosons. In the critical (“ten dimensional”) case, this is achieved by a chiral GSO projection [1]. In this section we will see that a straightforward generalization exists in the non-critical case as well, giving rise to a large class of stable string theories with the number of degrees of freedom varying between that of two dimensional string theory and that of the critical superstring. We will discuss the “type II” theory – as usual there exists a heterotic version.

Let us first recall how the GSO projection works in the case of ten dimensional (perhaps compactified) string theory. One starts with a fermionic string vacuum, which is normally left-right symmetric (non chiral GSO projection), and therefore contains two sectors (NS, NS) and (R, R) (for the left and right movers). Both sectors contain bosonic excitations in space-time – all states contribute with positive sign to the partition sum. As discussed above, such theories are tachyonic due to the large number of states ($c_{\text{string}} > 1$). A chiral GSO projection can be implemented if the theory has a chiral Z_2 R – symmetry. If there are no global anomalies, we can gauge this Z_2 symmetry (or in other words orbifold by it [1]); this involves two elements; first eliminating all states which are not invariant under the Z_2 symmetry from the spectrum. Hence only a subset of the original bosonic (NS, NS) and (R, R) states survive the projection. Then, as standard for orbifolds, there are also twisted sectors, which in this context are the (NS, R) and (R, NS) sectors, which contain space-time fermions. The theory thus obtained is not necessarily space-time supersymmetric, however under favorable circumstances it is tachyon free.

It is not known in general which Z_2 symmetries can be gauged and furthermore give rise to tachyon free “superstring” models. There is, however, a large class of theories, where this Z_2 is part of a larger chiral algebra, which are in general tachyon free and possess some additional nice properties. These are the space-time supersymmetric string vacua. In general such vacua can be constructed iff the original fermionic string vacuum has a global $N = 2$ superconformal symmetry². The projection can then be performed by constructing the (chiral) space-time SUSY charge S [28], and projecting out all operators which are not local w.r.t. S ; then one should again add “twisted sectors” obtained by acting with S on the remaining (NS, NS) and (R, R) operators; this gives rise to the (R, NS) and (NS, R) sectors (space-time fermions). This procedure is expected to be in general

² To avoid confusion it is important to emphasize that this $N = 2$ symmetry is an “accidental” global symmetry of the vacuum. It is not gauged (the BRST charge is that of the $N = 1$ fermionic string), and most excitations are not invariant under this symmetry – it is a property of the vacuum and not of the full theory. The existence of the $N = 2$ symmetry is a necessary and sufficient condition for space-time SUSY [27].

free of global anomalies (modular invariant), although no general proof exists. Also, such vacua are automatically free of tachyons, assuming there is only one time direction and the remaining “matter” is unitary. This follows from the space-time SUSY algebra. We will not elaborate on this procedure here, but rather explain directly how it generalizes to the non-critical case. For details on the “critical” construction we refer the reader to the original literature [28], [27], [29].

The procedure we are going to describe turns (non-critical) fermionic string vacua with an accidental global $N = 2$ symmetry to consistent superstrings. We will describe it in the particular case of the $D = d + 1$ dimensional fermionic string. The general structure is an obvious generalization, and can be found in [19].

The $d + 1$ dimensional fermionic string is described in superconformal gauge by d matter superfields X_i , $i = 1, \dots, d$, and the super Liouville field Φ . In components we have:

$$\begin{aligned} X_i &= x_i + \theta\psi_i + \bar{\theta}\bar{\psi}_i + \theta\bar{\theta}F_i; \quad i = 1, \dots, d \\ \Phi &= \phi_l + \theta\psi_l + \bar{\theta}\bar{\psi}_l + \theta\bar{\theta}F_l \end{aligned} \tag{3.1}$$

The fields X_i are free, while Φ is described by the super Liouville Largangian [11]:

$$S_{SL} = \frac{1}{2\pi} \int d^2z \int d^2\theta [D\Phi\bar{D}\Phi + 2\mu \exp(\alpha_+\Phi)] \tag{3.2}$$

where $D = \partial_\theta + \theta\partial_z$, and we have dropped curvature couplings (1.2). The central charge is $\hat{c}_L (= \frac{2}{3}c_L) = 1 + 2Q^2$, $Q = \sqrt{\frac{9-d}{2}}$. As mentioned above, this theory has a tachyonic ground state (for $d > 1$). The corresponding vertex operator is

$$T_k = \int d^2\theta \exp(ik \cdot X + \beta\Phi) \tag{3.3}$$

where $\frac{1}{2}k^2 - \frac{1}{2}\beta(\beta + Q) = \frac{1}{2}$. As in the bosonic case, $E = \beta + \frac{Q}{2}$ (see section 1 (1.6)), $E^2 - k^2 = m^2 = \frac{1-d}{8}$, so that T_k is a tachyon for $d > 1$.

For $d = 9$ (critical ten dimensional string theory), the Z_2 R – symmetry that one orbifolds by to obtain the superstring is $\psi \rightarrow -\psi$, $\bar{\psi} \rightarrow \bar{\psi}$ (or equivalently $\theta \rightarrow -\theta$, $\bar{\theta} \rightarrow \bar{\theta}$). The tachyon (3.3) transforms to minus itself under this symmetry, and is projected out of the spectrum. It is tempting to try and divide by the same symmetry for all d . Unfortunately, this is too naive; for $d < 9$ the orbifoldized theory doesn’t make sense; there are a number of ways of seeing that – modular invariance breaks down, the spin field [28] doesn’t have the necessary properties, etc. We have to use the more sophisticated approach related to global $N = 2$ symmetry. The basic observation is that for odd $d = 2n + 1$, such

that the total dimension of space-time is even³, the system (3.1) possesses an $N = 2$ symmetry. This symmetry pairs $d - 1 = 2n$ of the X_i into $N = 2$ superfields, and the remaining $X_{2n+1} \equiv X$ is paired with Φ . This involves breaking the $O(d)$ symmetry; one can think of Φ, X as light cone coordinates, while the rest of the X_i ($i = 1, \dots, 2n$) are transverse directions. The vacuum is not translationally invariant in the Φ direction. The action of the $N = 2$ superconformal symmetry on the transverse X_i is completely standard and we will not review it. The free Φ, X system forms a chiral $N = 2$ multiplet: it is convenient to define $\phi = \phi_l + ix$, $\psi = \psi_L + i\psi_x$, etc. The $N = 2$ generators are:

$$\begin{aligned}
T(z) &= -\frac{1}{2}\partial\phi\partial\phi^\dagger + \frac{1}{4}(\psi\partial\psi^\dagger + \psi^\dagger\partial\psi) - \frac{1}{4\gamma}\partial^2\phi - \frac{1}{4\gamma}\partial^2\phi^\dagger \\
J(z) &= \frac{1}{2}\psi\psi^\dagger - \frac{1}{2\gamma}\partial\phi + \frac{1}{2\gamma}\partial\phi^\dagger \\
T_F^+ &= \frac{1}{2}\psi^\dagger\partial\phi + \frac{1}{2\gamma}\partial\psi^\dagger \\
T_F^- &= \frac{1}{2}\psi\partial\phi^\dagger + \frac{1}{2\gamma}\partial\psi
\end{aligned} \tag{3.4}$$

In (3.4), $\dagger$ does not interchange left and right movers – it corresponds to complex conjugation in field space: $\phi^\dagger = \phi_l - ix$, etc. As in Liouville theory, the free field expressions (3.4) should be understood literally away from an interacting region (“wall”), which we haven’t specified yet. The string coupling is again (as in the bosonic theories) $g_{\text{st}} \propto e^{-\frac{Q}{2}\phi}$, and we need a potential to suppress the region $\phi \rightarrow -\infty$; we haven’t yet determined the appropriate potential. The form (3.4) is still very useful, for example to discuss bulk effects, which are anyway the best understood part of the Liouville correlation functions.

To choose an appropriate wall, it is convenient to define the chiral $N = 2$ superfield $\Gamma = \phi + \theta\psi + \bar{\theta}\bar{\psi} + \theta\bar{\theta}F + \dots$ (such that $D^\dagger\Gamma = 0$). Here θ is a complex Grassmanian variable (different from the θ in (3.1) – (3.3)). In terms of Γ , the free Lagrangian for Φ, X is:

$$\mathcal{L} = \frac{1}{8\pi} \int d^4\theta \Gamma \Gamma^\dagger$$

We would like now an $N = 2$ invariant operator to provide a potential for ϕ and to set the scale. There is a very natural choice for the potential, which preserves the $N = 2$ symmetry:

$$\mathcal{L} = \frac{1}{8\pi} \int d^4\theta \Gamma^\dagger \Gamma + i \frac{\mu}{8\pi\gamma} \int d^2\theta e^{-\gamma\Gamma} + i \frac{\mu}{8\pi\gamma} \int d^2\theta^\dagger e^{-\gamma\Gamma^\dagger} \tag{3.5}$$

³ There is an analogous requirement in compactified critical string theory.

The potential is an F-term, while from the $N = 1$ point of view it is a tachyon condensate at some particular non zero momentum. This value of the momentum is special because (3.5) is actually manifestly $N = 2$ supersymmetric; in fact, it is precisely the $N = 2$ Liouville Lagrangian. However its role here is to supply an accidental global symmetry – it is *not* a remnant of $N = 2$ supergravity in superconformal gauge. The parameter γ is not renormalized in $N = 2$ Liouville (unlike in $N = 0, 1$ Liouville): $\gamma = \frac{1}{Q}$. The $N = 2$ Liouville system (3.5) together with the $2n$ X_i forms a vacuum of ($N = 1$) fermionic string theory with a global $N = 2$ symmetry. However, this is not the superstring yet – the theory still contains tachyons (3.3), does not contain space-time fermions, etc. To perform the chiral GSO projection, we define the target space SUSY operator S :

$$S(z) = e^{-\frac{1}{2}\sigma + \frac{i}{2}(H_1 + \dots + H_n + H_l + Qx)} \quad (3.6)$$

Here σ is the bosonized superconformal ghost current [28]: $\beta\gamma = \partial\sigma$; the H_i are bosonized fermions: $\psi_{2i-1}\psi_{2i} = \partial H_i$ ($i = 1, \dots, n$), and similarly $\psi_l\psi_x = \partial H_l$. The GSO projection amounts to keeping in the spectrum only operators which have a local OPE with $S(z)$. Note that apart from the last term ($e^{\frac{i}{2}Qx}$) in S , this just means projecting by $(-)^{F_R}$, where F_R is the right moving fermion number (the naive projection $\psi \rightarrow -\psi$). For $Q = 0$ (the critical case) we recover the original GSO projection. In general, the projection ties the momentum in the x direction with the fermion number. In particular, p_x takes discrete values. The operator $S(z)$ is chiral ($\bar{\partial}S = 0$); this seems strange in view of the fact that x can be non-compact (so that only operators with $p_{\text{left}} = p_{\text{right}}$ are physical). The point is that the spectrum of x momenta is determined by locality w.r.t. S ; after the projection, the radius of x is not a meaningful concept.

After choosing the “longitudinal” direction, the theory still has $SO(2n)$ rotation invariance in the X_i directions. This means that the supercharge S (3.6) is not unique. Additional charges may be obtained by acting on it with the $SO(2n)$ generators $e^{\pm iH_a \pm iH_b}$, ($a \neq b = 1, 2, \dots, n$). This gives a set of charges S_α in one of the spinor representations of $SO(2n)$. To get the other representation (for $n > 0$), one uses the fact that the operator:

$$\tilde{S}(z) = e^{-\frac{1}{2}\sigma + \frac{i}{2}(-H_1 - \dots - H_{n-1} + H_n - H_l - Qx)} \quad (3.7)$$

is local w.r.t. S , and therefore is physical. Applying $SO(2n)$ rotations to $\tilde{S}$ generates the second spinor representation. For odd n the two are isomorphic, while for even n , S_α and $S_{\dot{\beta}}$ transform as different representations. The zero modes:

$$Q_\alpha = \oint \frac{dz}{2\pi i} S_\alpha(z); \quad \tilde{Q}_{\dot{\beta}} = \oint \frac{dz}{2\pi i} \tilde{S}_{\dot{\beta}}(z) \quad (3.8)$$

satisfy the super-algebra

$$\begin{aligned} \{Q_\alpha, \tilde{Q}_{\dot{\beta}}\} &= \gamma_{\alpha\dot{\beta}}^i P_i && \text{for } n \text{ even} \\ \{Q_\alpha, \tilde{Q}_\beta\} &= \gamma_{\alpha\beta}^i P_i && \text{for } n \text{ odd} \end{aligned} \quad (3.9)$$

The algebra (3.9) is a “space SUSY” algebra in the transverse directions: lightcone (ϕ_l, x) translations do not appear on the r.h.s., essentially because the vacuum is not translationally invariant in ϕ .

We now turn to discuss some general features of the spectrum of these theories. Although most states are paired by the SUSY algebra (3.9), apriori some tachyonic states could remain, since lack of tachyons is no longer implied by the algebra. However, for unitary matter theories (such as (3.1)), one can check directly in general that the theory is tachyon free after the GSO projection. From the point of view of section 2 this is clear: the effective density of states of the GSO projected theory (2.7) receives contributions only from states which are killed by the supercharges:

$$Q_\alpha |\text{phys}\rangle = \tilde{Q}_{\dot{\beta}} |\text{phys}\rangle = 0 \quad (3.10)$$

The rest of the states are paired level by level and cancel in the free energy (2.7). Now (3.10) implies that $|\text{phys}\rangle$ is independent of the transverse excitations (more generally of the transverse excitations and the internal degrees of freedom); the number of states that can satisfy (3.10) is therefore of the order of the number of states in the Φ, X (or Γ) system, which is a 2D string system. Hence $\lim_{s \rightarrow 0} G(s)$ is finite, and as shown in section 2, tachyons can not exist by modular invariance. Unitarity enters in two places:

- 1) Condition (2) of the theorem proved in section 2 breaks down in general in non unitary theories.
- 2) If the conditions are satisfied, the argument above proves only that the one loop partition sum is finite. There could be cancellations between NS and R tachyons, leading to a finite partition sum in the presence of tachyons. In unitary theories, Ramond tachyons can not occur (even before the chiral projection). Hence, only (NS, NS) tachyons may exist, and these are ruled out by the above argument.

Absence of tachyons can also be established directly [19], but we feel that the above argument is more intuitive. In the D dimensional superstring the explicit check of absence of tachyons is particularly simple: the only tachyon before the projection is T_k (3.3). The GSO projection (locality of T_k w.r.t. S (3.6)) implies: $k_x Q \in 2Z + 1$. The smallest value of

k_x (corresponding to the lightest state) is $|k_x| = \frac{1}{Q}$, for which $(\beta + \frac{Q}{2})^2 = \frac{Q^2}{4} + \frac{1}{Q^2} - 1 \geq 0$ for all Q . Hence the momentum in the x direction is quantized precisely such that the tachyon is always massive. In particular, the zero momentum tachyon (the original cosmological term in (3.2)) is projected out⁴.

Another interesting property of these theories is that target SUSY can be broken continuously, by switching on the modulus $\lambda \int \partial x \bar{\partial} x$. This perturbation is not projected out by GSO; it is clear from the form (3.6) of the generators that the symmetry is broken for any $\lambda \neq 0$. The situation is quite different from the ten dimensional case, where continuous breaking of SUSY can not occur [27]. The difference is due to the time dependence of the vacuum solution, but the detailed mechanism is still unclear; the gravitino is massive here even when SUSY is unbroken.

Returning to the $N = 2$ Liouville action, one can ask what is the role of the $N = 2$ cosmological term: why did we choose an $N = 2$ invariant perturbation to set the scale, out of the infinite number of GSO invariant operators, which are not $N = 2$ invariant (in general). After all, the analysis of the spectrum and symmetries are in any case performed in the region where the interaction is negligible. The answer is that it is indeed admissible to set the scale with other on shell physical vertex operators. Nevertheless, we do expect the $N = 2$ invariant perturbation (3.5) to exhibit especially nice properties. First, since (3.5) is $N = 2$ invariant, we have manifest space SUSY for all μ . For other perturbations in (3.5), SUSY would in general be broken. In addition, F-terms in $(2, 2)$ theories are known to yield moduli of the appropriate CFT (see e.g. [30]). For generic perturbations, the dynamics of the corresponding “Liouville” model (3.5) is probably much more complicated.

The $N = 2$ cosmological constant has $\beta = -\gamma = -\frac{1}{Q}$, or $E = \beta + \frac{Q}{2} = \frac{Q}{2} - \frac{1}{Q}$. This is positive only when $Q^2 > 2$. As $Q^2 \rightarrow 2$ ($d \rightarrow 5$), the lowest lying NS state goes to zero mass, and for $d > 5$, its energy becomes negative (while the mass increases). As mentioned above, the behavior of $E > 0$ and $E < 0$ states is qualitatively different. One would need a more detailed understanding of the theory to see what changes as $E \rightarrow 0$, but it is clear that the point $Q^2 = 2$ corresponds to some kind of transition in the behavior of the theory. For $Q^2 < 2$ there are other possibilities to set the scale in (3.5) with operators of positive energy, however they are all less symmetric. The special role of this new “critical dimension” is not entirely clear, and should be elucidated further.

⁴ Note also that the remaining modes of the tachyon go to infinite mass as $Q \rightarrow 0$. Therefore, they are invisible in the critical theory.

The GSO procedure may superficially resemble fine tuning the coefficients of all the tachyonic modes in the action to zero. The difference between this and fine tuning is that in the GSO procedure the states are projected out by gauge invariance (the Z_2 R symmetry). Therefore, we are assured that they will not reappear in various intermediate channels in higher loops. On the other hand, as we saw in section 2, fine tuning does not solve the problem of higher loops (topologies).

An interesting property of the non-critical superstrings we have constructed is that they have a vanishing partition sum (generically) to all orders in the genus expansion. Consider first the partition sum on the sphere: usually, the spherical partition sum is not zero in non-critical string theory [31]. The reason is the following. In critical string theory, the partition sum vanishes due to the six $c, \bar{c}$ (reparametrization ghost) zero modes on the sphere. However, in the non-critical case, the Liouville path integral contains a comparable divergence, due to zero modes of the classical Liouville solution. The zero related to the $c, \bar{c}$ zero modes is really $\frac{1}{\text{vol}SL(2,C)}$, and the infinity in the Liouville sector is proportional to the same volume (up to a finite factor), thus the two cancel. More precisely, choosing the conformal gauge does not fix the gauge completely. By fixing the $SL(2, C)$ invariance and therefore not integrating over the various zero modes, one finds a finite answer. In fermionic string theory a similar phenomenon occurs, with $SL(2, C)$ replaced by $OSp(2, 1)$. Again, the infinite factors cancel, and one is left with a finite partition sum. In the limit $Q \rightarrow 0$ (when the non-critical string approaches a critical one), we still have a finite partition sum, but exactly at $Q = 0$, ϕ translation invariance appears, and we have to divide this finite answer by the volume of ϕ . The free energy per unit volume $\frac{\text{const}}{V} \rightarrow 0$ as $V \rightarrow \infty$ (ϕ is non compact). Thus, in space-time the difference between the critical and the non-critical cases is that in the latter, due to lack of ϕ translation invariance, we compute the total free energy, which is finite, while in the critical case, where translational invariance is restored, we are interested in the free energy per unit volume, which is zero.

When the fermionic string vacuum possesses a global $N = 2$ symmetry (even before the chiral projection, which is of course irrelevant for the partition sum on the sphere), there are *two additional* Liouville fermion zero modes, obtained by applying $N = 2$ transformations to the usual fermionic zero modes, which exist for $N = 1$ Liouville. The ghosts, which are still those of the $N = 1$ string do not have balancing zero modes, therefore the path integral vanishes. This means that the classical vacuum energy vanishes in (3.5). The torus, and higher genus partition sums also vanish for $n \geq 1$, by the usual contour deformation arguments [28], [32]. In that case, bosons and fermions are paired, except perhaps for

a set of measure zero of states at zero momentum $p_i = 0$ ($i = 1, \dots, 2n$). Sometimes, such arguments can be subtle due to contact terms from boundaries of moduli space, however here we expect such subtleties to be absent in general, since there are no massless excitations.

To illustrate the above abstract discussion, we finish this section with a brief analysis of the simplest theories constructed here: the two and four dimensional superstrings ($n = 0, 1$).

Example 1: Two dimensional superstring⁵.

This theory contains two superfields: the super Liouville field Φ , and a space coordinate X ; the two combine into an $N = 2$ Liouville system (3.4) with $\gamma(= \frac{1}{Q}) = \frac{1}{2}$. To calculate the torus partition sum, we have to sum (2.1) over all states satisfying:

- (a) Locality w.r.t. (3.6); here, $S(z) = e^{-\frac{1}{2}\sigma + \frac{i}{2}H + iX}$.
- (b) $\Delta - \bar{\Delta} \in Z$.
- (c) Mutual locality.

There are four sectors in the theory. We will next go over them and solve the conditions (a) – (c).

Consider first the (NS, NS) sector: the vertex operators have the form

$$e^{-\sigma + inH_l + ipX} e^{-\bar{\sigma} + i\bar{n}\bar{H}_l + i\bar{p}\bar{X}}$$

Condition (a) leads to

$$\begin{aligned} p &= \frac{n-1}{2} + m \\ \bar{p} &= \frac{\bar{n}-1}{2} + \bar{m} \end{aligned} \tag{3.11}$$

($m, \bar{m} \in Z$). From condition (b) we have

$$\begin{aligned} n - \bar{n} &\in 2Z \\ p^2 - \bar{p}^2 &\in 2Z \end{aligned} \tag{3.12}$$

while from condition (c):

$$pp' - \bar{p}\bar{p}' \in Z \tag{3.13}$$

for all p, p' in the (NS, NS) spectrum. The solution of the constraints (3.11) – (3.13) is:

- (1) $n, \bar{n} \in 2Z + 1$; $p = m$, $\bar{p} = \bar{m}$, $m - \bar{m} \in 2Z$.

⁵ Based on [33].

(2) $n, \bar{n} \in 2Z$; $p = m + \frac{1}{2}$, $\bar{p} = \bar{m} + \frac{1}{2}$, $m - \bar{m} \in 2Z$.

Summing (3.4) over these states gives (after multiplying by the oscillator contribution for X , H_l , and the ghost contribution):

$$Z_{\text{NS,NS}} = \frac{\sqrt{\tau_2}}{2} |\theta_2|^2 \quad (3.14)$$

A similar analysis applied to the other 3 sectors gives

$$Z_{\text{NS,R}} = Z_{\text{R,NS}} = -\frac{\sqrt{\tau_2}}{4} |\theta_2|^2 \quad (3.15)$$

$$Z_{\text{R,R}} = \sqrt{\tau_2} [|\theta_2|^2 + |\theta_3|^2 + |\theta_4|^2] \quad (3.16)$$

To see better where the different contributions are coming from, we should list the physical operators which survive the projection. Chirally, in the NS sector we have the tachyon operators:

$$T_k = e^{-\sigma + ikX + \beta\phi_l}; \quad k \in Z + \frac{1}{2} \quad (3.17)$$

The restriction on the spectrum of k 's is due to the requirement of locality with $S(Z)$ (3.6).

In the Ramond sector we have two sets of operators of the form:

$$V_k = e^{-\frac{1}{2}\sigma + \frac{i}{2}\epsilon H + ikX + \beta\phi_l} \quad (3.18)$$

with

$$\begin{aligned} (1) \quad & \epsilon = 1 \quad k = n = 0, 1, 2, 3, \dots; \quad \beta = n - 1 \\ (2) \quad & \epsilon = -1 \quad k = -l - \frac{1}{2} \quad l = 0, 1, 2, 3, \dots; \quad \beta = l - \frac{1}{2} \end{aligned} \quad (3.19)$$

In agreement with the partition sums (3.14) – (3.16). Due to the low space-time dimension, the superalgebra (3.9) is quite degenerate: there is only one supercharge, $Q = \oint \frac{dz}{2\pi i} S(z)$, (3.6), which satisfies $Q^2 = 0$. Hence, Q is a BRST like charge. An interesting theory is obtained by restricting ourselves to $\ker Q$: physical operators are those which satisfy:

$$Q|\text{phys}\rangle = 0 \quad (3.20)$$

This gives a topological string theory (reminiscent of the topological model of the bosonic string with $c = -2$ matter [34]). The condition (3.20) projects out the tachyon modes T_k (3.17) with $k < 0$, and the second set of Ramond states in (3.19) (those with $\epsilon = -1$, $k < 0$). Since the physical states must have the same Liouville momentum for left

and right movers (ϕ_l is non compact), all space-time fermions are projected out by (3.20). The topological two dimensional superstring contains (NS, NS) tachyons

$$T_n = e^{-\sigma + i(n + \frac{1}{2})x + (n - \frac{1}{2})\phi_l}; \quad n = 0, 1, 2, \dots \quad (3.21)$$

and (R, R) states:

$$V_n = e^{-\frac{1}{2}\sigma + \frac{i}{2}H_l + inx + (n-1)\phi_l}; \quad n = 0, 1, 2, \dots \quad (3.22)$$

(here we mean left-right symmetric combinations $\bar{V}_{\text{left}} V_{\text{right}}$). Correlation functions of the operators (3.21), (3.22) can be obtained using the methods of [35]. We leave further investigation of this topological theory to future work.

If one does not impose (3.20) on the spectrum, one can ask how does the theory change when we add the (physical) operator $\partial x \bar{\partial} x$ to the action. The dependence of the partition sum on the “radius” of x can be obtained by viewing the theory as a chiral orbifold. The idea is the following: if x has radius R ($x \simeq x + 2\pi R$), we can study the orbifold theory obtained by gauging the Z_2 symmetry:

$$\begin{aligned} x &\rightarrow x + \pi R \\ H_l &\rightarrow H_l + \pi \\ \bar{H}_l &\rightarrow \bar{H}_l \\ \sigma &\rightarrow \sigma - i\pi \\ \bar{\sigma} &\rightarrow \bar{\sigma} + 2\pi i \end{aligned} \quad (3.23)$$

This formulation of the theory as a chiral orbifold emphasizes the role of the Z_2 symmetry which is being gauged (3.23). The role of the space SUSY operator $S(z)$ is obscure from this point of view. In particular, the symmetry generator Q (3.8) is in the spectrum only for particular R ($R = 2$). This symmetric point is not singled out in (3.23).

By standard methods one obtains the partition sums:

$$Z_{NS,NS}(R) = \sqrt{\tau_2} \sum_{m,m' \in \mathbb{Z}} q^{\frac{1}{2}(\frac{2m+1}{R} + \frac{m'}{2})^2} \bar{q}^{\frac{1}{2}(\frac{2m+1}{R} - \frac{m'}{2})^2} \quad (3.24)$$

$$Z_{R,R}(R) = \sqrt{\tau_2} \sum_{m,m' \in \mathbb{Z}} q^{\frac{1}{2}(\frac{m}{R} + \frac{m'}{2})^2} \bar{q}^{\frac{1}{2}(\frac{m}{R} - \frac{m'}{2})^2} \quad (3.25)$$

$$Z_{NS,R}(R) + Z_{R,NS}(R) = -\sqrt{\tau_2} \sum_{m,m' \in \mathbb{Z}} q^{\frac{1}{2}(\frac{2m}{R} + \frac{R}{2}(m' + \frac{1}{2}))^2} \bar{q}^{\frac{1}{2}(\frac{2m}{R} - \frac{R}{2}(m' + \frac{1}{2}))^2} \quad (3.26)$$

For $R = 2$ we have the results (3.14), (3.15), (3.16). For generic R , the total partition sum can be written as a linear combination of $c = 1$ partition sums: $Z_{\text{total}}(R) = 2Z_{c=1}(R) - Z_{c=1}(\frac{4}{R})$. The integral $\Omega = \int_{\mathcal{F}} \frac{d^2\tau}{\tau^2} Z_{\text{total}}$, which as we have learnt in section two gives the number of degrees of freedom, can be performed using (2.11) (or the results of [36]); one finds that $\Omega \propto R$. In particular, at the “supersymmetric”, or topological, point $R = 2$, the cancellation of bosons and fermions is not complete. This is due to the low space-time dimension – the SUSY algebra $Q^2 = 0$ does not imply pairing of bosons and fermions. As explained above, we need at least two non compact transverse directions for that. Therefore, we will next consider the case of two transverse dimensions.

Example 2: Four dimensional superstring.

The “matter” consists of two superfields X_1, X_2 (in addition to Φ, X), $Q = \sqrt{3}$, and $S = e^{-\frac{1}{2}\sigma + \frac{i}{2}(H_l + H_1 + \sqrt{3}x)}$. One can repeat the discussion of Example 1 here; we will leave the details to the reader, and only give the final result for the one loop partition sums⁶:

$$\begin{aligned}
\sqrt{\tau_2}|\eta(\tau)|^6 Z_{NS,NS} = & |\theta_3 + \theta_4|^2 \sum_{\epsilon} \Theta \left[\frac{\frac{\epsilon+3}{6}}{0} \right] (12\tau) \bar{\Theta} \left[\frac{\frac{\epsilon+3}{6}}{0} \right] (12\bar{\tau}) + \\
& |\theta_3 - \theta_4|^2 \sum_{\epsilon} \Theta \left[\frac{\frac{\epsilon}{6}}{0} \right] (12\tau) \bar{\Theta} \left[\frac{\frac{\epsilon}{6}}{0} \right] (12\bar{\tau}) + \\
& (\theta_3 + \theta_4)(\bar{\theta}_3 - \bar{\theta}_4) \sum_{\epsilon} \Theta \left[\frac{\frac{\epsilon+3}{6}}{0} \right] (12\tau) \bar{\Theta} \left[\frac{\frac{\epsilon}{6}}{0} \right] (12\bar{\tau}) + \\
& (\theta_3 - \theta_4)(\bar{\theta}_3 + \bar{\theta}_4) \sum_{\epsilon} \Theta \left[\frac{\frac{\epsilon}{6}}{0} \right] (12\tau) \bar{\Theta} \left[\frac{\frac{\epsilon+3}{6}}{0} \right] (12\bar{\tau})
\end{aligned} \tag{3.27}$$

$$\begin{aligned}
\sqrt{\tau_2}|\eta(\tau)|^6 Z_{R,R} = & |\theta_2|^2 \sum_{\epsilon} \Theta \left[\frac{\frac{\epsilon-\frac{1}{2}}{6}}{0} \right] (12\tau) \bar{\Theta} \left[\frac{\frac{\epsilon-\frac{1}{2}}{6}}{0} \right] (12\bar{\tau}) + \\
& |\theta_2|^2 \sum_{\epsilon} \Theta \left[\frac{\frac{\epsilon+3-\frac{1}{2}}{6}}{0} \right] (12\tau) \bar{\Theta} \left[\frac{\frac{\epsilon+3-\frac{1}{2}}{6}}{0} \right] (12\bar{\tau}) + \\
& |\theta_2|^2 \sum_{\epsilon} \Theta \left[\frac{\frac{\epsilon-\frac{1}{2}}{6}}{0} \right] (12\tau) \bar{\Theta} \left[\frac{\frac{\epsilon+3-\frac{1}{2}}{6}}{0} \right] (12\bar{\tau}) + \\
& |\theta_2|^2 \sum_{\epsilon} \Theta \left[\frac{\frac{\epsilon+3-\frac{1}{2}}{6}}{0} \right] (12\tau) \bar{\Theta} \left[\frac{\frac{\epsilon-\frac{1}{2}}{6}}{0} \right] (12\bar{\tau}) +
\end{aligned} \tag{3.28}$$

⁶ All sums over ϵ run over $\epsilon = 0, 2, 4$.

$$\begin{aligned}
-\sqrt{\tau_2}|\eta(\tau)|^6 Z_{NS,R} = & (\theta_3 + \theta_4)\bar{\theta}_2 \sum_{\epsilon} \Theta \left[\frac{\epsilon+3}{6} \right] (12\tau) \bar{\Theta} \left[\frac{\epsilon+\frac{3}{2}}{6} \right] (12\bar{\tau}) + \\
& (\theta_3 - \theta_4)\bar{\theta}_2 \sum_{\epsilon} \Theta \left[\frac{\epsilon}{6} \right] (12\tau) \bar{\Theta} \left[\frac{\epsilon-\frac{3}{2}}{6} \right] (12\bar{\tau}) + \\
& (\theta_3 + \theta_4)\bar{\theta}_2 \sum_{\epsilon} \Theta \left[\frac{\epsilon+3}{6} \right] (12\tau) \bar{\Theta} \left[\frac{\epsilon-\frac{3}{2}}{6} \right] (12\bar{\tau}) + \\
& (\theta_3 - \theta_4)\bar{\theta}_2 \sum_{\epsilon} \Theta \left[\frac{\epsilon}{6} \right] (12\tau) \bar{\Theta} \left[\frac{\epsilon+\frac{3}{2}}{6} \right] (12\bar{\tau}) +
\end{aligned} \tag{3.29}$$

Now, by general arguments above, we expect $Z_{\text{total}} = Z_{NS,NS} + Z_{R,R} + Z_{NS,R} + Z_{R,NS}$ to vanish! Quite miraculously from the point of view of the explicit formulae (3.27) – (3.29), one can write Z_{total} in a much simpler way as:

$$\sqrt{\tau_2}|\eta(\tau)|^6 Z_{\text{total}} = \sum_{\epsilon} |F_{\epsilon}(\tau)|^2 \tag{3.30}$$

where

$$\begin{aligned}
F_{\epsilon}(\tau) = & (\theta_3 + \theta_4)(\tau) \Theta \left[\frac{\epsilon+3}{6} \right] (12\tau) + (\theta_3 - \theta_4)(\tau) \Theta \left[\frac{\epsilon}{6} \right] (12\tau) \\
& - \theta_2(\tau) \left\{ \Theta \left[\frac{\epsilon+\frac{3}{2}}{6} \right] (12\tau) + \Theta \left[\frac{\epsilon-\frac{3}{2}}{6} \right] (12\tau) \right\}
\end{aligned} \tag{3.31}$$

The most natural scenario would be to have $F_{\epsilon}(\tau) = 0$ for all $\epsilon (= 0, 2, 4)$. This seems indeed to be the case. We have checked the vanishing of F_{ϵ} (3.31) for low orders in q , and believe that the result is true in general, although we haven't proved it analytically. If this is the case, we have, as expected, $Z_{\text{total}} = 0$.

4. CLASSICAL DYNAMICS IN 2D STRING THEORY.

In the previous sections we have seen that stable (tachyon free) string vacua are in general those with $c_{\text{string}} = 1$. Bosonic strings with $c_{\text{string}} = 1$ (two dimensional strings), are known from the matrix models [5], [6], [37], [38] to be solvable. In this section we will review the understanding of their properties in the continuum path integral formalism, in the hope that some of them may carry over and help to understand more complicated theories with $c_{\text{string}} = 1$, like those of section 3.

In the introduction, we have seen that the set of “resonant” amplitudes defines for all d a “bulk” S – matrix, which is given by Shapiro – Virasoro type integral representations.

To illustrate this, we discuss here tachyon dynamics in $D = d+1$ dimensional “non-critical” string theory. Of course, for generic D , there is no reason to concentrate on the tachyon, both because it is merely the lowest lying state of the string spectrum, and because it is tachyonic, thus absent in more physical theories (see section 3). Our justification is that for $D = 2$ it is the only field theoretic degree of freedom, and is massless.

The on shell vertex operator for the tachyon is:

$$T_k = \exp(ik \cdot X + \beta(k)\phi) \quad (4.1)$$

where k, X are d – vectors, and by (1.5):

$$\frac{1}{2}k^2 - \frac{1}{2}\beta(\beta + Q) = 1 \quad (4.2)$$

which implies as before (1.8): $m_{\text{tachyon}}^2 = \frac{2-D}{12}$. The bulk amplitudes described in the introduction take here the form:

$$A_{s=0}(k_1, ..k_N) = \langle T_{k_1} .. T_{k_N} \rangle / \log \mu \quad (4.3)$$

where $\sum_i k_i = 0$, and by (1.9) $\sum_i \beta_i = -Q$. As mentioned above, such amplitudes have an integral representation for all D :

$$A_{s=0}(k_1, .., k_N) = \prod_{i=4}^N \int d^2 z_i |z_i|^{2(k_1 \cdot k_i - \beta_1 \beta_i)} |1 - z_i|^{2(k_3 \cdot k_i - \beta_3 \beta_i)} \prod_{4=i < j}^N |z_i - z_j|^{2(k_i \cdot k_j - \beta_i \beta_j)} \quad (4.4)$$

For $D = 26$ this is the familiar Shapiro – Virasoro representation [1], however it retains all the nice properties for all D . The singularities, which are the most important features of (4.4), are poles in various channels, corresponding to on shell intermediate particles. To examine these poles, one focuses on contributions to (4.4) of regions in moduli space where vertices approach each other. E.g. from the region $z_4, z_5, ..., z_{n+2} \rightarrow 0$, we have an infinite series of poles in (E, p) , $E = \frac{Q}{2} + \sum_i \beta_i$, $p = \sum_i k_i$ ($i = 1, 4, 5, 6, ..., n+2$), with residues related to lower correlation functions with insertions of various intermediate states. For example, near the first (tachyon) pole we have:

$$A_{s=0}(k_1, .., k_N) \simeq \frac{\langle T_{k_1} T_{k_4} ... T_{k_{n+2}} T_{\bar{k}} \rangle \langle T_{\sum_i k_i} T_{k_2} T_{k_3} T_{k_{n+3}} ... T_{k_N} \rangle}{(\frac{Q}{2} + \sum_i \beta_i)^2 - (\sum_i k_i)^2 - \frac{2-D}{12}} \quad (4.5)$$

where $\tilde{k} = -\sum_i k_i$, and similarly for the higher poles. Thus, for generic correlators (4.4), the pole structure is very complicated. For $N = 4$, a closed expression for (4.4) in terms of Γ functions is known [1]:

$$A_{s=0}(k_1, \dots, k_4) = \pi \prod_{i=2}^4 \frac{\Gamma(k_1 \cdot k_i - \beta_1 \beta_i + 1)}{\Gamma(\beta_1 \beta_i - k_1 \cdot k_i)} \quad (4.6)$$

It is easy to check that (4.6) satisfies factorization (4.5). For $N \geq 5$ no such closed forms are known.

The situation is markedly different in two dimensions. There, the singularity structure of (4.4) is much simpler, and the integral representation can be evaluated for all N . The condition (4.2) implies that we have left and right moving massless “tachyons” in space-time:

$$T_k^{(\pm)} = \exp(ikX + \beta^{(\pm)}\phi); \quad \beta^{(\pm)} = -\sqrt{2} \pm k \quad (4.7)$$

The correlation functions have then the following form [39]:

$$\begin{aligned} \langle T_{k_1}^{(+)} \dots T_{k_{N-1}}^{(+)} T_{k_N}^{(-)} \rangle &= (N-3)! \prod_{i=1}^N (-\pi) \frac{\Gamma(\frac{1}{2}\beta_i^2 - \frac{1}{2}k_i^2)}{\Gamma(1 - \frac{1}{2}\beta_i^2 + \frac{1}{2}k_i^2)} \\ \langle T_{k_1}^{(+)} \dots T_{k_n}^{(+)} T_{p_1}^{(-)} \dots T_{p_m}^{(-)} \rangle &= 0; \quad n, m \geq 2 \end{aligned} \quad (4.8)$$

The Γ factor for $i = N$ in (4.8) is infinite. This infinity should be interpreted as the volume of the Liouville mode $\log \mu$ (see section 1). The vanishing of the second set of correlators in (4.8) can be understood as a lack of such a volume factor. Taking this into account, the first form properly understood describes all bulk N point functions [39], [35].

The results (4.8) are at first sight very puzzling. Both the vanishing of amplitudes with $n, m \geq 2$, and the simple form of the amplitudes with $m = 1$ (or $n = 1$) are not at all obvious from (4.4); apriori one would expect *much more* poles, as in (4.5). We will not derive (4.8) here (the reader may find a more complete discussion in [39], [35]), but we will explain the origin of its simple form. The phenomenon behind this simplicity is partial decoupling of a certain infinite set of discrete states. To understand why discrete states are important, we have to recall the spectrum of the two dimensional string [40], [22], [41]. We have already encountered the tachyon field (4.7). For $D > 2$ there is in addition an infinite tower of massive transverse oscillator states. For $D = 2$ there are no transverse directions, hence most of the massive degrees of freedom are absent. There are still oscillator states at discrete values of the momentum, which are related to oscillator

primary states of $c = 1$ CFT [42] (in space-time these are longitudinal oscillations of the string). The general form of these states is:

$$V_{r_1, r_2}^{(\pm)} = [\partial^{r_1 r_2} X + \dots] \exp \left(i \frac{r_1 - r_2}{\sqrt{2}} X + \beta_{r_1, r_2}^{(\pm)} \phi \right) \quad (4.9)$$

where $r_1, r_2 \in Z_+$, and the $\dots$ stands for a certain polynomial in $\partial X, \partial^2 X, \dots$ (with scaling dimension $r_1 r_2$). The Liouville dressing is $\beta_{r_1, r_2}^{(\pm)} = -\sqrt{2} \pm \frac{r_1 + r_2}{\sqrt{2}}$; note that unlike (4.7), here $V^{(+)}$ ($V^{(-)}$) correspond to positive (negative) energy states.

The states (4.9) together with the tachyon comprise the full spectrum of 2d string theory (at ghost number zero). Thus, all N point functions (4.4) must factorize on these states. To demonstrate the role of the discrete states in (4.4), we should analyze the factorization in various channels. Consider, for example, the simplest degeneration channel, when two tachyons collide. There are two possibilities (up to $X \rightarrow -X$): 1) $T_{k_1}^{(+)}(z)T_{k_2}^{(-)}(0)$; 2) $T_{k_1}^{(+)}(z)T_{k_2}^{(+)}(0)$, $z \rightarrow 0$.

In the first case, using the free OPE, which is clearly valid in bulk correlators, although not in general [8], [20], we find an infinite set of poles at $k_1 k_2 + (k_1 - \sqrt{2})(k_2 + \sqrt{2}) = -n$, $n = 1, 2, 3, \dots$. The first pole ($n = 1$) corresponds to an intermediate tachyon, while the rest must have vanishing residues. The reason is that the residue involves correlators of oscillator states (generalizing (4.5)), but the momentum $k_1 + k_2$ of the intermediate state is *not* in the discrete list (4.9) (at least for generic k_1, k_2), therefore the corresponding state must be BRST trivial (null), and decouple.

In the second case, the poles occur at $k_1 k_2 - (k_1 - \sqrt{2})(k_2 - \sqrt{2}) = -n$ or: $\sqrt{2}(k_1 + k_2) = 2 - n$. The intermediate momentum is automatically in the discrete list (4.9); furthermore $\beta = \beta_1 + \beta_2 = -\sqrt{2} - \frac{n}{\sqrt{2}}$ is such that the energy of the intermediate state is negative – it is a $V^{(-)}$. A similar situation occurs for poles in more complicated channels.

Therefore, the pole structure of (4.4) depends crucially on the physics of the states $V_{r_1, r_2}^{(-)}$. All but a finite number of the poles in the tachyon S – matrix correspond to these states. One can show [35] that they decouple from amplitudes of tachyons,

$$\langle V_{r_1, r_2}^{(-)} T_{k_1} \dots T_{k_N} \rangle = 0 \quad (4.10)$$

as long as $k_1, \dots, k_N \notin \frac{1}{\sqrt{2}}Z$. Therefore, most of the poles in (4.4) have vanishing residues. Using (4.10) and factorization it is straightforward to show that most of the tachyon amplitudes vanish, and the ones with signatures $(N - 1, 1)$ have the simple structure (4.8) (see [35]).

The poles in (4.8) appear as “external leg factors” [22], however they also have a standard interpretation as poles in the (i, N) channels (i.e. when T_{k_i} and T_{k_N} approach each other) – the only channels where the poles have finite residues. Since k_N is fixed by kinematics ($k_N = -\frac{N-2}{\sqrt{2}}$), they appear to be “leg poles”. One could ask at this point why these poles have a finite residue; after all, the intermediate states are again $V^{(-)}$ in this case. The point is that the decoupling (4.10) of the discrete states with negative energies is *only partial*. If there are enough $V^{(+)}$ ’s, the amplitude does not vanish. The residues of the poles in the (i, N) channel involve the three point function $\langle V_{r_1, r_2}^{(-)} T_{k_1} T_{k_2} \rangle$. By the conservation laws, k_1, k_2 are in this case in the discrete list (4.9) and their energies are positive; this correlation function does not vanish. The general situation is the following: the most general correlator has the form:

$$\langle V_{r_1, r_2}^{(+)} \dots V_{r_{2n-1}, r_{2n}}^{(+)} V_{s_1, s_2}^{(-)} \dots V_{s_{2l-1}, s_{2l}}^{(-)} T_{k_1} \dots T_{k_N} \rangle \propto (\log \mu)^{n-l} \quad (4.11)$$

where $k_1, \dots, k_N \notin \frac{1}{\sqrt{2}}\mathbb{Z}$. Eq. (4.11) is very useful to determine the relative size of various amplitudes. If the power of $\log \mu$ ($n-l$) is one, we have a finite bulk amplitude. This is the case for (4.3) with signature $(N-1, 1)$ and generic $k_1, \dots, k_{N-1}$; k_N is fixed by kinematics to be $V_{0, N-2}^{(+)}$. If the power $n-l \leq 0$, the bulk amplitude (coefficient of $\log \mu$) vanishes. This is the case for tachyon amplitudes with signature (n, m) , $n, m \geq 2$, and generic momenta (here the conservation laws do not constrain individual momenta), and for correlation functions like (4.10).

One natural question that arises at this point is whether we can reconstruct the general (non bulk) amplitudes in the theory from the knowledge of the bulk ones. In general, this issue is not understood (although it is probably possible to do this). However, in 2d (closed) string theory it appears that one can make an educated guess for the answer, based on a physical assumption. Since the results obtained by this method are in agreement with those obtained from matrix models, the physical assumption is probably justified, however it has not been derived from first principles (in the continuum formalism). The idea is the following: we saw before how the structure of 2d string theory leads to the bulk correlators (4.8). It is very natural to define “renormalized operators” ($\Delta(x) \equiv \frac{\Gamma(x)}{\Gamma(1-x)}$):

$$\tilde{T}_k = \frac{T_k}{(-\pi)\Delta(\frac{1}{2}\beta^2 - \frac{1}{2}k^2)} \quad (4.12)$$

These have very simple correlation functions:

$$\langle \tilde{T}_{k_1} \dots \tilde{T}_{k_N} \rangle = (N-3)! \quad (4.13)$$

What happens for non zero s (1.9)? Integer $s > 0$ is easy to treat since if $\langle \tilde{T}_{k_1} \dots T_{k_N} \rangle = \tilde{A}_{k_1 \dots k_N} \mu^s$, then differentiating s times w.r.t. μ we find

$$\tilde{A}(k_1, \dots, k_N) = \frac{\langle \tilde{T}_{k_1} \dots \tilde{T}_{k_N} (\tilde{T}_{k=0})^s \rangle}{s!}$$

The numerator is a bulk amplitude, so we can use (4.13) and find: $A(k_1, \dots, k_N) = \frac{(s+N-3)!}{s!} \mu^s$, or:

$$A_s(k_1, \dots, k_N) = \left(\frac{\partial}{\partial \mu} \right)^{N-3} \mu^{s+N-3} \quad (4.14)$$

We can further redefine the operators (and rescale the path integral) such that all μ dependence on the r.h.s. of (4.14) disappears; alternatively put $\mu = 1$. Then we find ourselves in a situation where all correlators at integer s are polynomials in $k_1, \dots, k_N$ (using (1.9)). The question is how we can continue to non integer s . The first guess would be to declare (4.14) valid for any s – it certainly makes sense for all amplitudes. While this is true in a certain region in momentum space, in general the answer is more subtle.

There are several clues pointing in the direction we should choose. The first clue is the fact that the effect of the massive modes of the string is summarized, at least for bulk amplitudes, in the external leg factors (4.8). The renormalized field $\tilde{T}_k$ (4.12) is completely insensitive to the massive modes. This suggests that the S – matrix of $\tilde{T}$ is described by a local two dimensional field theory. One might have expected to see tachyon poles in amplitudes, however as we saw, they are not there (4.8). The reason is lack of conservation of Liouville momentum in this field theory (1.9). From the point of view of Liouville theory, we have the OPE:

$$\exp(\beta_1 \phi) \exp(\beta_2 \phi) = \int d\beta \exp(\beta \phi) f(\beta, \beta_1, \beta_2) \quad (4.15)$$

where f is an OPE coefficient. The contour of integration for β should [8] run over macroscopic states $\beta = -\frac{Q}{2} + ip$ ($p \in R$). Using the OPE (4.15) in generic tachyon correlation functions we find that the contribution of regions of moduli space where vertices approach each other is:

$$\begin{aligned} \int d^2 z \langle T_{k_1}(z) T_{k_2}(0) \dots \rangle &\simeq \int_{|z| < \epsilon} d^2 z \int_{-\infty}^{\infty} dp (z \bar{z})^{\frac{1}{2} p^2 + \frac{1}{2} (k_1 + k_2)^2 - 1} f(p, \beta_1, \beta_2) \langle T_{k_1 + k_2} \dots \rangle \\ &\simeq \int_{-\infty}^{\infty} dp \frac{f(p, \beta_1, \beta_2)}{p^2 + (k_1 + k_2)^2} \langle T_{k_1 + k_2} \dots \rangle \end{aligned} \quad (4.16)$$

where we interchanged the order of the z and p integrations. For fixed p , (4.16) has the familiar form from critical string theory; we find a pole corresponding to the intermediate state $T_{k_1+k_2}$. The fact that Liouville momentum is not conserved and we have to sum over all p 's may turn this pole into a cut: (4.16) depends on $|k_1 + k_2|$. Thus we expect cuts whenever some of the momenta $\{k_i\}$ in (4.3) satisfy $\sum_i k_i \equiv p \rightarrow 0$. The fact that the bulk $\tilde{T}$ correlators are polynomial in k_i (4.14) is another indication that $\tilde{T}$ is described by a local 2d field theory. However, due to the expected appearance of cuts in amplitudes, we have to be careful in continuing (4.14) from its region of validity. The remaining issue is to understand the expected analytic structure of the correlation functions.

First, if the picture we have been developing is correct, we expect the three point functions $\langle T_{k_1} T_{k_2} T_{k_3} \rangle$ to be given exactly by (4.14):

$$\langle \tilde{T}_{k_1} \tilde{T}_{k_2} \tilde{T}_{k_3} \rangle = 1 \quad (4.17)$$

The reason is that non analytic effects such as (4.16) can only occur for $N \geq 4$ point functions – three point functions are insensitive to the non-conservation of energy (4.15). This immediately allows us to obtain the propagator of the tachyon. By putting $k_2 = 0$ in (4.17) and integrating we obtain the two point function⁷:

$$\langle \tilde{T}_k \tilde{T}_{-k} \rangle = \frac{1}{\sqrt{2}|k|} \quad (4.18)$$

Hence the propagator (in a convenient normalization) should be: $\frac{|k|}{\sqrt{2}}$. An important consistency check on this is the comparison of an amplitude with an insertion of a puncture $P = T_{k=0}$ to the amplitude without it. By KPZ scaling (1.9) we have:

$$\langle P T_{k_1} \dots T_{k_N} \rangle = \left[\frac{1}{\sqrt{2}} \sum_{i=1}^N |k_i| - (N - 2) \right] \langle T_{k_1} \dots T_{k_N} \rangle \quad (4.19)$$

Thinking of (4.19) as a relation between tree amplitudes in the purported space time field theory reveals its essential features: we can insert the puncture $T_{k=0}$ into the tree amplitude $\langle T_{k_1} \dots T_{k_N} \rangle$ either by attaching it to one of the N external legs, thus adding an internal propagator of momentum $k_i + 0 = k_i$ or inside the diagram. The first term

⁷ In critical string theory the two point function vanishes, however, as explained above for the partition sum, this is due to a division by the (infinite) volume of ϕ . Here, due to lack of translational invariance, the two point function is finite, and is related to the propagator.

(the sum) on the r.h.s. of (4.19) corresponds to the first possibility; we can read off the propagator $\frac{|k|}{\sqrt{2}}$, which agrees with that obtained from (4.18). The second term corresponds to the second possibility. Now, after understanding the propagator, the remaining problem is specifying the vertices in the space-time field theory. The three point vertex is 1 (4.17).

The higher vertices can be calculated using two basic properties:

a) The form (4.14) is exact for $k_1, \dots, k_{N-1} > 0$, $k_N < 0$, which is the region in which all the energies involved are positive (4.3); from the world sheet point of view this is natural since positive energy perturbations correspond to small deformations of the surface [8], and also because the integrals (4.4) converge there (after a certain analytic continuation in the central charge [35]).

b) All higher irreducible vertices are *analytic* in $\{k_i\}$. To understand this, consider for example the four point function:

$$\tilde{A}(k_1, \dots, k_4) = \int d^2z \langle \tilde{T}_{k_1}(0) \tilde{T}_{k_2}(\infty) \tilde{T}_{k_3}(1) \tilde{T}_{k_4}(z) \rangle \quad (4.20)$$

We can separate the z integral in (4.20) into two pieces. One is a sum of three contributions from the regions $z \rightarrow 0, 1, \infty$. By (4.16) we expect to get from those the tachyon propagator⁸ $\frac{1}{\sqrt{2}}|k_1 + k_i|$. The rest of the z integral is the contribution of the bulk of moduli space (where z is not close to $0, 1, \infty$); it gives a new irreducible four particle interaction (which we will denote by $A_{1PI}^{(4)}$) for the tachyons. This contribution is analytic, since only massless intermediate states cause cuts in the amplitudes (4.16), and we have subtracted their contribution. Using this decomposition, we expect the following analytic structure for $\tilde{A}(k_1, \dots, k_4)$:

$$\tilde{A}(k_1, \dots, k_4) = \left(\frac{|k_1 + k_2|}{\sqrt{2}} + \frac{|k_1 + k_3|}{\sqrt{2}} + \frac{|k_1 + k_4|}{\sqrt{2}} \right) + A_{1PI}^{(4)} \quad (4.21)$$

where $A_{1PI}^{(4)}$ is analytic in k_i . Next we use the fact that in the region $k_1, k_2, k_3 > 0$, $k_4 < 0$, we actually know $\tilde{A}$ (4.14). Comparing to (4.21) we find

$$A_{1PI}^{(4)} = -1 \quad (4.22)$$

Eq. (4.22) holds now everywhere in $\{k_i\}$ by analytic continuation.

⁸ The contribution of the massive states in the OPE (4.16) is presumably related to the factors $\Delta(\frac{1}{2}\beta_i^2 - \frac{1}{2}k_i^2)$ in (4.12).

It is now clear how to proceed in the case of N point functions. We assume that we know already $A_{1PI}^{(4)}, \dots, A_{1PI}^{(N-1)}$. Then we write all possible tree graphs with N external legs, propagator $\frac{|k|}{\sqrt{2}}$ and vertices $A_{1PI}^{(n)}$ ($n \leq N-1$) and add an unknown new irreducible vertex $A_{1PI}^{(N)}(k_1, \dots, k_N)$. $A_{1PI}^{(N)}$ is again analytic in $\{k_i\}$ and we can fix it by comparing the sum of exchange amplitudes (reducible graphs) and $A_{1PI}^{(N)}$ to the full answer (4.14) in the appropriate kinematic region. This fixes $A_{1PI}^{(N)}$ in the above kinematic region. Now we use analyticity of $A_{1PI}^{(N)}$ to fix it everywhere. The outcome of this process is the determination of the amplitudes in all kinematic regions given their values in one kinematic region.

In practice, it is more convenient to obtain $A_{1PI}^{(N)}$ by Legendre transforming. This gives a highly non trivial set of irreducible vertices; the general formulae are quite involved [35]. As an example, one finds:

$$A_{1PI}^{(N)}(k_1, k_2, k_3, k_4 = 0, \dots, k_N = 0) = (\partial_\mu)^{N-3} \left\{ \frac{1}{\mu} \prod_{i=1}^3 \frac{1}{\cosh(\frac{k_i}{\sqrt{2}} \log \mu)} \right\} \Big|_{\mu=1} \quad (4.23)$$

$$A_{1PI}^{(N)}(k_1, k_2, k_3, k_4, 0, \dots, 0) = (\partial_\mu)^{N-4} \mu^{-2} \left(\prod_{i=1}^4 \frac{1}{\cosh(\frac{k_i}{\sqrt{2}} \log \mu)} \right) \left(-1 - \mu \partial_\mu \log \prod_{1 \leq i < j \leq 3} \cosh(\frac{k_i + k_j}{\sqrt{2}} \log \mu) \right) \Big|_{\mu=1} \quad (4.24)$$

As expected, the irreducible vertices are analytic in $\{k_i\}$. This is true in general. Note that the preceding discussion is very reminiscent of the decomposition of moduli space which appears in general in closed string field theory [43]. It would be interesting to make the relation more precise. The tachyon field theory constructed above is also closely related to the one which arises in matrix models [44], [45], [46]).

This concludes our presentation of tree level scattering in 2d closed string theory. While we now know the structure of the S – matrix in great detail, our understanding of the origin of the factorization of poles (4.8) and the emergence of the local tachyon field theory is still unsatisfactory. For example, does the spectrum (massless tachyon + discrete states) automatically lead to the structure we have described? In particular, is the decoupling of the discrete states with $E < 0$ generic?

To gain more information about the possible structures in 2d string theory, we have studied other string models with similar general characteristics. The first is the 2d fermionic string. For $N = 1$ supergravity [35], the space-time dynamics includes two massless scalars and a set of discrete states. The classical S – matrix is extremely similar

to the bosonic case: the discrete states with $E \leq 0$ again (partially) decouple, and the bulk S – matrix again has only factorized poles, as in (4.8).

The 2d $N = 2$ (critical) string was studied in [47], and its tree level S – matrix is even simpler (all $N \geq 4$ point functions vanish). This is due to the fact that the theory is really four dimensional. Open 2d strings exhibit quite different behavior from the closed case [48]. The spectrum contains again the massless field (4.7) and discrete states (4.9), however the S – matrix of the open string tachyon $T_k^{(B)}$ on the disk has the form ($m_i = \frac{1}{2}\beta_i^2 - \frac{1}{2}k_i^2$):

$$A_{\text{open}}^{(n,m)} = \left[\prod_{i=1}^{n+m} \frac{1}{\Gamma(1 - m_i)} \right] H_n(k_1, \dots, k_n) H_m(p_1, \dots, p_m) \quad (4.25)$$

where

$$H_n(k_1, k_2, \dots, k_n) = \sum_{\sigma} \prod_{l=1}^{n-1} \frac{1}{\sin \pi \sum_{i=1}^l m_{\sigma(i)}} \quad (4.26)$$

and the sum over σ in (4.26) runs over permutations of $1, 2, \dots, n$. The form (4.25), (4.26) exhibits in general a rich pole structure which is in essence due to the fact that the states $V_{r_1, r_2}^{(-)}$ *do not* decouple here. Therefore, the decoupling of the massive states from the tachyon dynamics also doesn't occur here. Remarkably, the complicated S – matrix (4.25), (4.26) again follows from a very simple space-time field theory. Indeed, one can check [48] that the generating functional:

$$\mathcal{W}(T^{(\pm)}) = \int D\psi D\psi^* D\bar{\psi} D\bar{\psi}^* \exp \left[- \int dX d\phi e^{\sqrt{2}\phi} \mathcal{L}(\psi, T) \right] \quad (4.27)$$

where the Lagrangian $\mathcal{L}$ is given by:

$$\mathcal{L}(\psi, T) = \psi^* \sin(\pi\sqrt{2}\partial^+) \psi + \bar{\psi}^* \sin(\pi\sqrt{2}\partial^-) \bar{\psi} + T^{(+)} \bar{\psi}^* \bar{\psi} + T^{(-)} \psi^* \psi, \quad (4.28)$$

and $\partial^{\pm} = \partial_{\phi} \pm \partial_X$. satisfies:

$$\frac{\delta^{n+m} \mathcal{W}}{\delta T_{k_1}^{(+)} \dots \delta T_{k_n}^{(+)} \delta T_{p_1}^{(-)} \dots \delta T_{p_m}^{(-)}} = H_n(k_1, \dots, k_n) H_m(p_1, \dots, p_m) \quad (4.29)$$

The reader should consult [48] for details and a more precise formulation. Hence, (4.27) generates the scattering amplitudes (4.25), (4.26). Furthermore, the propagator in (4.28) is essentially a lattice propagator, and it is natural to define the theory on a space-time lattice in $\frac{1}{2}(X \pm i\phi_l)$. Thus, the structure arising in the open sector is both richer (4.25), (4.26) and simpler (4.27), (4.28) than the one observed in the closed sector of the theory.

5. COMMENTS.

Despite the recent progress, there are still many unresolved fundamental questions both in 2d string theory, and in its relation to higher dimensional models. We have identified one feature which 2d string theory shares with higher dimensional stable theories – the fact that $c_{\text{string}} = 1$ (the small effective number of states). It is not clear what other features of 2d string theory have higher dimensional analogs. The non critical superstring models of section 3 may be useful to study this issue.

Even within the framework of 2d string theory, the understanding of the free fermion structure of [5], [6] is very incomplete. This seems to boil down to a better understanding of the dynamics of $V_{r,s}^{(\pm)}$ (4.9). Understanding the dynamics of the discrete states is crucial also to study gravitational physics in 2d string theory. The black hole of [25] is described [49] by turning on $V_{1,1}^{(-)}$; other $V_{r,s}^{(-)}$ seem also to play a role in gravitational back reaction.

Two dimensional closed string theory seems to be described by two consistent (but different) S – matrices. The S – matrix for T_k (4.1), given by (4.4) is sensitive to the discrete states (4.9), and contains the information on gravitational physics. Its pole structure and the effective action which describes it may be used to study the space-time 2d gravity. On the other hand, $\tilde{T}_k$ (4.12) are described by a second S – matrix, which originates from a two dimensional field theory for the tachyon field. In particular, it doesn't contain space-time gravity. The appearance of the local tachyon field theory is quite surprising from the point of view of the continuum formulation (as is the case with many other features of these models, it arises more naturally in the matrix model [44], [45], [46]); it would be interesting to understand this dual structure better.

A very important question concerns the form of the exact classical equations of motion in 2d string theory. The correlation functions we have found suggest a very interesting structure. In the sigma model approach, one can investigate the moduli space of classical solutions by requiring that adding perturbations to the world action: $\mathcal{S} \rightarrow \mathcal{S} + \lambda_{\Delta} \int V_{\Delta}$, does not spoil conformal invariance [50] [51]. From the form of the correlation functions of section 4 and matrix models [37], [38] it follows that adding $\lambda_k \int T_k$ to the action for all $k \notin \frac{1}{\sqrt{2}}Z$ and any λ_k doesn't spoil conformal invariance. We can calculate all correlators in a power series in λ_k (for fixed μ), and find sensible results. This seems to suggest that the exact non linear classical equations of motion of 2d string theory have the property that a tachyon field of the form $T(X) = \sum_{k \notin \frac{1}{\sqrt{2}}Z} T_k$ (and trivial metric and other fields) is a solution. Of course, this can only be true up to field redefinitions, but even then it is

quite remarkable (a special case of this is the claim that Liouville (1.2) is a CFT). When we turn on an expectation value of a tachyon with one of the discrete momenta, the metric and other fields back react. The details of this back reaction haven't been worked out yet.

The higher genus correlation functions are another issue, which is still not resolved in the continuum formalism. In the matrix model approach, all order correlation functions were obtained in [37]. This is especially interesting in the light of the simple results obtained; as an example, the two point function was found to be given by:

$$\frac{\partial}{\partial \mu} \langle T_k T_{-k} \rangle = \frac{\Gamma(-\sqrt{2}|k|)}{\Gamma(\sqrt{2}|k|)} \text{Im} \left\{ e^{\frac{i\pi}{\sqrt{2}}|k|} \left[\frac{\Gamma(\frac{1}{2} + \sqrt{2}|k| - i\mu)}{\Gamma(\frac{1}{2} - i\mu)} - \frac{\Gamma(\frac{1}{2} - i\mu)}{\Gamma(-\sqrt{2}|k| + \frac{1}{2} - i\mu)} \right] \right\} \quad (5.1)$$

Higher point functions appear in [37]. To extract the result for given genus one should expand (5.1) in powers of $\frac{1}{\mu^2}$. It is challenging to derive the higher genus correlation functions in the continuum formalism, and even more challenging to understand the origin of the “non-perturbative” results, such as (5.1).

Acknowledgements

I would like to thank N. Seiberg for numerous discussions of the issues presented here; part of the work described in this review was done in collaboration with him, and the rest benefitted greatly from his suggestions and criticism. I would also like to thank M. Bershadsky and P. Di Francesco for collaboration and discussions. I have benefitted from discussions with T. Banks, R. Dijkgraaf, J. Distler, P. Freund, D. Gross, I. Klebanov, E. Martinec, G. Moore, N. Seiberg, S. Shatashvili, S. Shenker and H. Verlinde. I am grateful to the organizers of the spring school and ICTP for their hospitality. This work was partially supported by DOE grant DE-AC02-76ER-03072.

References

- [1] M. Green, J. Schwarz and E. Witten, *Superstring theory*, Cambridge Univ. Press (1987) and references therein.
- [2] A. Polyakov, *Gauge Fields and Strings*, Harwood Academic Publishers (1987).
- [3] E. Brezin, C. Itzykson, G. Parisi and J.B. Zuber, Comm. Math. Phys. **59** (1978) 35 ; D. Bessis, Comm. Math. Phys. **69** (1979) 147
- [4] V. Kazakov, Phys. Lett. **150B** (1985) 282; F. David, Nucl. Phys. **B257** (1985) 45; J. Ambjorn, B. Durhuus and J. Frohlich, Nucl. Phys. **B257** (1985) 433; V. Kazakov, I. Kostov and A. Migdal, Phys. Lett. **157B** (1985) 295.
- [5] D. Gross and A. Migdal, Phys. Rev. Lett. **64** (1990) 127; M. Douglas and S. Shenker, Nucl. Phys. **B335** (1990) 635; E. Brezin and V. Kazakov, Phys. Lett. **B236** (1990) 144.
- [6] E. Brezin, V. Kazakov, and Al. Zamolodchikov, Nucl. Phys. **B338** (1990) 673; D. Gross and N. Miljković, Phys. Lett **238B** (1990) 217; P. Ginsparg and J. Zinn-Justin, Phys. Lett **240B** (1990) 333; G. Parisi, Phys. Lett. **238B** (1990) 209.
- [7] T. Banks, M. Douglas, N. Seiberg and S. Shenker, Phys. Lett. **238B** (1990) 279.
- [8] N. Seiberg, Rutgers preprint RU-90-29 (1990).
- [9] G. Moore, N. Seiberg and M. Staudacher, Rutgers preprint RU-91-11 (1991).
- [10] G. Moore and N. Seiberg, Rutgers preprint RU-91-29 (1991).
- [11] A. Polyakov, Phys. Lett. **B103** (1981) 207, 211.
- [12] V. Knizhnik, A. Polyakov and A. Zamolodchikov, Mod. Phys. Lett **A3** (1988) 819.
- [13] F. David, Mod. Phys. Lett. **A3** (1988) 1651; J. Distler and H. Kawai, Nucl. Phys. **B321** (1989) 509.
- [14] S. Das, S. Naik and S. Wadia, Mod. Phys. Lett. **A4** (1989) 1033; J. Polchinski, Nucl. Phys. **B324** (1989) 123; T. Banks and J. Lykken, Nucl. Phys. **B331** (1990) 173.
- [15] A. Belavin, A. Polyakov and A. Zamolodchikov, Nucl. Phys. **B241**, 333 (1984).
- [16] T. Curtright and C. Thorn, Phys. Rev. Lett. **48** (1982) 1309; E. Braaten, T. Curtright and C. Thorn, Phys. Lett. **118B** (1982) 115; Ann. Phys. **147** (1983) 365.
- [17] J.L Gervais and A. Neveu, Nucl. Phys. **B209** (1982) 125; **B224** (1983) 329; **B238** (1984) 125.
- [18] D. Kutasov and N. Seiberg, Nucl. Phys. **B358** (1991) 600.
- [19] D. Kutasov and N. Seiberg, Phys. Lett. **B251** (1990) 67.
- [20] J. Polchinski, Texas preprints UTTG-19-90; UTTG-39-90 (1990).
- [21] A. Gupta, S. Trivedi and M. Wise, Nucl. Phys. **B340** (1990) 475.
- [22] A. Polyakov, Mod. Phys. Lett. **A6** (1991) 635.
- [23] A. B. Zamolodchikov JETP Lett. **43** (1986) 730; J. Cardy and A. Ludwig, Nucl. Phys. **B285**[**FS19**] (1987) 687; A. Cappelli, D. Friedan and J. Latorre, Rutgers preprint RU-90-43 (1990).

- [24] E. Witten, IAS preprint IASSNS-HEP-91/51 (1991).
- [25] E. Witten, Phys. Rev. **D44** (1991) 314.
- [26] J. Polchinski, Comm. Math. Phys. **104** (1986) 37; B. Maclain and B. Roth, Comm. Math. Phys. **111** (1987) 539; K. O' Brien and C. Tan, Phys. Rev. **D36** (1987) 1184.
- [27] D. Friedan, A. Kent, S. Shenker and E. Witten, unpublished; T. Banks, L. Dixon, D. Friedan and E. Martinec, Nucl. Phys. **B299** (1988) 613.
- [28] D. Friedan, E. Martinec and S. Shenker, Nucl. Phys. **B271** (1986) 93.
- [29] D. Gepner, Nucl. Phys. **B296** (1988) 757; PUPT-1121 (1989).
- [30] L. Dixon, Princeton preprint PUPT-1074 (1987).
- [31] A.B. Zamolodchikov, Phys. Lett. **117B** (1982) 87.
- [32] E. Martinec, Phys. Lett. **171B** (1986) 189.
- [33] D. Kutasov, G. Moore and N. Seiberg, unpublished.
- [34] E. Witten, Nucl. Phys. **B340** (1990) 281; J. Distler, Nucl. Phys. **B342** (1990) 523.
- [35] P. Di Francesco and D. Kutasov, Princeton preprint PUPT-1276 (1991).
- [36] M. Bershadsky and I. Klebanov, Phys. Rev. Lett. **65** (1990) 3088; N. Sakai and Y. Tanii, Tokyo preprint TIT/HEP-160 (1990).
- [37] G. Moore, Rutgers preprint RU-91-12 (1991).
- [38] D. Gross and I. Klebanov, Princeton preprint PUPT-1241 (1991); G. Mandal, A. Sengupta, and S. Wadia, IAS preprint, IASSNS-HEP/91/8 (1991); K. Demeterfi, A. Jevicki, and J.P. Rodrigues, Brown preprints BROWN-HET-795, 803 (1991); J. Polchinski, Texas preprint UTTG-06-91 (1991).
- [39] P. Di Francesco and D. Kutasov, Phys. Lett. **261B** (1991) 385.
- [40] N. Seiberg, unpublished.
- [41] B. Lian and G. Zuckerman, Yale preprint YCTP-P18-91 (1991); S. Mukherji, S. Mukhi and A. Sen, Tata preprint TIFR/TH/91-25 (1991); P. Bouwknegt, J. McCarthy and K. Pilch, preprint CERN-TH.6162/91 (1991).
- [42] J. Goldstone, unpublished; V. Kac, in *Group Theoretical Methods in Physics*, Lecture Notes in Physics, vol. 94 (1979); G. Segal, Comm. Math. Phys. **80** (1981) 301.
- [43] H. Sonoda and B. Zwiebach, Nucl. Phys. **B331** (1990) 592; **B336** (1990) 185.
- [44] S. Das and A. Jevicki, Mod. Phys. Lett. **A5** (1990) 1639.
- [45] A. Sengupta and S. Wadia, Tata preprint TIFR/TH/90-13 (1990).
- [46] D. Gross and I. Klebanov, Nucl. Phys. **B352** (1991) 671.
- [47] H. Ooguri and C. Vafa, Mod. Phys. Lett. **A5** (1990) 1389; Nucl. Phys. **B361** (1991) 469.
- [48] M. Bershadsky and D. Kutasov, Princeton preprint PUPT-1283 (1991).
- [49] M. Bershadsky and D. Kutasov, Phys. Lett. **266B** (1991) 345.
- [50] C. Callan, D. Friedan, E. Martinec and M. Perry, Nucl. Phys. **B262** (1985) 593.
- [51] T. Banks and E. Martinec, Nucl. Phys. **B294** (1987) 733.